

## PREVALENCE OF AFLATOXIN B<sub>1</sub> IN FINISHED COMMERCIAL BROILER FEED FROM WEST CENTRAL PAKISTAN

N. Rashid, M. A. Bajwa, M. Rafeeq, M. A. Khan, Z. Ahmad, M. M. Tariq, A. Wadood, and F. Abbas

Center for Advanced Studies in Vaccinology and Biotechnology, University of Balochistan, Quetta, Pakistan

Corresponding author E-mail: nadeemuob@yahoo.com

### ABSTRACT

Aflatoxin contamination of poultry feed results in heavy economic loss and a potential risk for human being. Finished commercial broiler feed (n=96) samples collected from different broiler farms for the period June 2009 to May 2010 from south west Pakistan were analyzed for aflatoxin B<sub>1</sub>(AfB<sub>1</sub>) by using thin layer chromatography with a detection limit of 3µg/kg. Analysis revealed that 91.66% samples were positive with a range 10 to 166µg/kg and mean 47.64±2.55µg/kg. Highest level of AfB<sub>1</sub> was found in the month of September ranged from 42 to 166µg/kg with mean of 83.25±16.11µg/kg. Categorically 82.30% samples were above the permissible limit recommended by United States Food and Drug Administration. A significant (P<0.05) effect of season on aflatoxin production was noted. The prevalence of AfB<sub>1</sub> may be attributed to poor harvesting technique, insect infestation, storage or environmental conditions.

**Key words:** aflatoxin, broiler, poultry feed, prevalence.

### INTRODUCTION

Aflatoxins are a group of structurally related toxic compounds produced predominantly by certain strains of *Aspergillus flavus*, and *Aspergillus parasiticus* (Diaz *et al.*, 2001). These molds are ubiquitous in nature and can grow and produce aflatoxins on any feed or grain used for poultry under favourable environmental conditions (Qazi and Fayyaz, 2006). United States (US) Food and Drug Administration (FDA) consider Aflatoxins to be an unavoidable contaminant of foods and have set regulatory levels. The permissible level for poultry is 20 ppb (Aravind *et al.*, 2003; Azab *et al.*, 2005). Aflatoxins are well intended the most prevalent, dangerous and studied mycotoxins (Krnjaja *et al.*, 2008). Ingestion of feed contaminated with aflatoxins causes high morbidity and mortality resulting severe economic losses (Oguz *et al.*, 2003; Diaz *et al.*, 2008). Aflatoxins not only impairs weight gain, feed intake, feed conversion efficiency, egg production (Ortatatli *et al.*, 2005), but also increases the susceptibility to environmental stress (Abdolamir *et al.*, 2005) and severity of diseases like crop mycosis, salmonellosis, coccidiosis, aspergillosis, and Marek's disease (Ibrahim *et al.*, 2000).

Although 18 different aflatoxins have been identified, only aflatoxins B<sub>1</sub>, B<sub>2</sub>, G<sub>1</sub>, and G<sub>2</sub> are significant contaminant of agricultural commodities and pose a potential risk to poultry (Shareef *et al.*, 2008). Among these, aflatoxin B<sub>1</sub> (AfB<sub>1</sub>) is of great concern due to its carcinogenic, mutagenic, teratogenic (Shi *et al.*, 2009), immunosuppressive (Hussain *et al.*, 2010) and growth inhibitory effects (Khan *et al.*, 2010). AfB<sub>1</sub> is

usually the major aflatoxin produced by toxigenic strains (Qazi and Fyyaz, 2006).

The incidence and level of Aflatoxin contamination in food and feeds is continuously monitored and many surveys on natural occurrence are conducted every year in industrialized countries (Diaz *et al.*, 2001). However, very little information is generated in developing countries. The inadequate technical, human and economic resources may be a reason for this lack of information (Cespedes and Diaz, 1997).

Keeping in view the importance of aflatoxins present study was planned to find out prevalence of aflatoxin B<sub>1</sub> in finished commercial broiler feed in and around Quetta district of Pakistan where no documented data is available. In addition to ascertain the prevalence of aflatoxin B<sub>1</sub>, the effect of season on its concentration and categorization of the level of contamination following the general guidelines of FDA in finished commercial broiler feed was also studied.

### MATERIALS AND METHODS

A base line study was conducted from June 2009 to May 2010 in and around Quetta district, situated between 30°15 North latitude and 66° 55 East longitudes, at an altitude of 1,675 meters in the North West part of Balochistan and at the West Central Pakistan. Quetta, is the capital city and the most populated district of Balochistan which is the largest province comprising the 45% of land mass of Pakistan. There are approximately 150 broiler farms with a flock rearing capacity of 1000 to 6000 birds.

### Prevalence of aflatoxin B<sub>1</sub>

**Sampling, Processing and Analysis:** The finished commercial broiler feed samples (n=96, 1kg each) were randomly collected on weekly basis from two government and private poultry farms in and around Quetta District. Grab method (Richard, 2000) was used to collect a fairly representative sample. The samples were collected in sterilized polythene bags and shifted to the toxicology laboratory of Center for Advanced Studies in Vaccinology and Biotechnology, University of Balochistan (CASVAB, UoB) within 24 hours for further processing and analysis.

The collected samples were dried at 80°C for 3 hours, grounded, passed from sieve No. 20 and stored in a double layer paper bag enveloped with polythene bag at -18°C until they were analyzed (within one week of collection).

Samples were analyzed quantitatively by Thin Layer Chromatography (TLC) to determine AfB<sub>1</sub> contents. Extraction and cleanup process were carried out by using the method no. 993.17, while quantification was done by using the method no. 998.03 of AOAC (2000).

**Effect of Season on AfB<sub>1</sub> concentration:** The effect of season on AfB<sub>1</sub> concentration was studied by dividing the whole year into four quarters viz., quarter 1 (June to August), quarter 2 (September to November), quarter 3 (December to February) and quarter 4 (March to May) as suggested by Anjum (1990).

**Categorization of level of contamination of AfB<sub>1</sub>** Collected samples were categorized into low (<20µg/kg), medium (20-50µg/kg) and high (>50µg/kg) level of contamination of AfB<sub>1</sub> following the general guidelines of FDA (Hanif *et al.*, 2006).

**Statistical Analysis:** Data revealed was subjected to statistical analysis using the One-Way Analysis of Variance as a Complete Randomized Design (CRD) using the general linear models procedure of statistical analysis software (SPSS 16 for windows). The means showing significant differences (P<0.05) were compared using Duncan's Multiple Range Test (DMR).

## RESULTS

Prevalence of AfB<sub>1</sub> in finished commercial broiler feed samples (n=96) is presented in Table 1. It was found that 91.6% of the examined samples were contaminated with AfB<sub>1</sub>. In the present study detection limit was 3µg/kg and the concentration of AfB<sub>1</sub> in positive samples ranged from 10 to 166µg/kg (mean 47.64±2.55µg/kg). The level of contamination was highest during the month of September (mean

83.25±16.11µg/kg) followed by October and August (mean 62.75±7.04 and 55.25±4.48µg/kg, respectively). The lowest level of contamination was found in the month of January that ranged from 10 to 30µg/kg with mean of 18.80±3.81 µg/kg. A significant difference in the means of AfB<sub>1</sub> was observed during months (P<0.05) Table 1. The effect of season on aflatoxin is depicted in Figure 1. The results revealed that highest concentration of AfB<sub>1</sub> was found in the quarter September to November (mean 66.04±7.03µg/kg), while it was lowest in the quarter December to February (mean 34.42±4.63µg/Kg). Following the general guidelines of FDA about contamination level of aflatoxins in poultry feed the results of the study reveals that 17.71% samples were categorized in low, 51.04% medium and 31.25% in high level of contamination (Table 2).


**Table 1. Spectrum of Aflatoxin B<sub>1</sub> in broiler feed with a detection limit of 3µg/kg**

Month	Total Samples	% +ve	Mean ± SEM for +ve samples (µg/Kg)	Range for +ve samples (µg/Kg)
Jun. 2009	8	87.5	40.42±7.05 <sup>bc</sup>	12-58
Jul. 2009	8	100	48.12±2.52 <sup>bc</sup>	39-58
Aug. 2009	8	100	55.25±4.48 <sup>bc</sup>	39-70
Sep. 2009	8	100	83.25±16.11 <sup>a</sup>	42-166
Oct. 2009	8	100	62.75±7.04 <sup>b</sup>	39-100
Nov. 2009	8	87.5	50.14±9.12 <sup>bc</sup>	17-83
Dec. 2009	8	100	55.37±3.35 <sup>bc</sup>	42-70
Jan. 2010	8	62.5	18.80±3.81 <sup>d</sup>	10-30
Feb. 2010	8	75	19.50±3.45 <sup>d</sup>	10-30
Mar. 2010	8	100	36.50±1.62 <sup>cd</sup>	30-42
Apr. 2010	8	87.5	41.71±1.82 <sup>bc</sup>	36-50
May 2010	8	100	40.75±1.58 <sup>bc</sup>	36-50
<b>Overall</b>	<b>96</b>	<b>91.66</b>	<b>47.64±2.55</b>	<b>10-166</b>

<sup>abcd</sup> Values within same column followed by different superscript differ significantly (P<0.05)

Table 2. Categorization of Aflatoxin B<sub>1</sub> contamination in poultry feed with respect to General guidelines of FDA

Month	Total samples	Level of Contamination (%)		
		Low contamination (<20 µg/Kg)	Medium contamination (20-50 µg/Kg)	High contamination (>50 µg/Kg)
Jun. 2009	8	37.5	37.5	25
Jul. 2009	8	0	75	25
Aug. 2009	8	0	37.5	62.5
Sep. 2009	8	0	25	75
Oct. 2009	8	0	25	75
Nov. 2009	8	25	25	50
Dec. 2009	8	0	37.5	62.5
Jan. 2010	8	75	25	0
Feb. 2010	8	75	25	0
Mar. 2010	8	0	100	0
Apr. 2010	8	12.5	87.5	0
May 2010	8	0	100	0
Overall	96	17.71	51.04	31.25

Fig. 1. Effect of Season on Prevalence of Aflatoxin B<sup>1</sup>

## DISCUSSION

Contamination of commercial feed with aflatoxins is not an unusual phenomenon. Several workers (Cespedes and Diaz, 1997; Charoenpornsook and Kavisarasai, 2006; Shareef, 2010 and Moalem *et al.*, 2010) from different parts of the world have reported incidence of aflatoxins form 40 to 92%. The maximum level of AfB<sub>1</sub> concentration observed in the current study was 166µg/kg. In other studies the highest values detected by Bhatti *et al.* (2001) Azab *et al.* (2005), Mngadi *et al.* (2008) and Alkhalaf *et al.* (2010) were 78, 120, 2000, and 70 µg/kg, respectively.

The differences in the prevalence and levels of contamination of aflatoxins in finished poultry feeds depend on number of factors such as the geographical area, climatic conditions (Dersjant-Li *et al.*, 2003), type of raw material used, birds and insects damage to grains used in formula feed (Thompson and Henke, 2000), and storage conditions (Moss, 2002). Similarly sampling procedure and preparation of samples is an important tool

that plays a vital role in the integrity of test result produced (Davis *et al.*, 1980).

The season has a major impact on fungal growth and aflatoxin production. The aflatoxin production is influenced by environmental conditions like humidity and temperature (Hanif *et al.*, 2008, Sultana and Hanif, 2009). In the present study high levels of AfB<sub>1</sub> in the quarter September to November and low concentrations during December to February indicates the effect of season. In Pakistan, monsoon (hot season with heavy rain fall in which there is persistent high relative humidity and temperature) commences from July to September (Yunus *et al.*, 2009) provide conditions ideal for the fungal growth ultimately resulting in increased production of aflatoxins.

According to the general guidelines of FDA about contamination level of aflatoxins in poultry feed the results of the present study are alarming that 82.30 % samples were above the permissible limits of 20ppb as recommended by USFDA (Ghahri, 2010). Such high level of aflatoxin in feed is not only hazardous to poultry because of direct exposure but also suggests a high risk to human health with the possibility of indirect exposure through contaminated meat, eggs and other poultry products and by products (Maqbool *et al.*, 2004; Bintvihok and Kositcharoenkul, 2006).

The present study clearly describes the existing status of aflatoxin B<sub>1</sub> contamination of finished commercial broiler feed in Quetta. This situation needs to be considered with strict surveillance and quality control of poultry feed.

## REFERENCES

Abdolamir, A., A. Safamehr, S.A. Mirhadi, M. Shivazad, M. Razzaghi-Abyaneh and A. Ashar-Naderi

- (2005). Evaluation of biochemical and production parameters of broiler chicks fed ammonia treated aflatoxin contaminated maize grains. *Animal Feed science and Technology*, 122: 289-301.
- Alkhalaf, N. A., A. K. Osman and K. A. Salama (2010). Monitoring of aflatoxins and heavy metals in some poultry feeds. *African Journal of Food Science*, 4: 192-199.
- Anjum A.D. (1990). Weather and disease: 1. Prevalence of the poultry disease in and around Faisalabad and their relationship to weather. *Pak. Vet. J.*, 10: 42-45.
- AOAC. (2000). *Natural Toxins. Official method of analysis*, 17<sup>th</sup> Edit., Association of Official Analytical Chemists. Arrington, Virginia. USA. : 11-12, 16-18.
- Aravind, K. L., V. S. Patil, G. Devegowda, B. Umakantha and S.P. Ganpule (2003). Efficacy of esterified glucomannan to counteract mycotoxicosis in naturally contaminated feed on performance and serum biochemical and hematological parameters in broilers. *Poultry Science*, 82: 571-576.
- Azab, R. M., W. M. Tawakkol, A. M. Hamad, M. K. Abou-Elmagd, H. M. El-Agrab and M. K. Refai (2005). Detection and estimation of aflatoxin B<sub>1</sub> in feeds and its biodegradation by bacteria and fungi. *Egyptian Journal of Natural Toxins*, 2: 39-56.
- Bhatti, B. M., T. Talat and R. Sardar (2001). Estimation of aflatoxin B<sub>1</sub> in feed ingredients and compound poultry feeds. *Pak. Vet. J.*, 21: 57-60.
- Bintvihok, A. and S. Kositchaenkul (2006). Effect of dietary calcium propionate on performance, hepatic enzyme activities and aflatoxin residues in broilers fed a diet containing low levels of aflatoxin B<sub>1</sub>. *Toxicon*, 47: 41-45.
- Cespedes, A. E. and G. J. Diaz (1997). Analysis of aflatoxins in poultry and pig feeds and feedstuffs used in Colombia. *Journal of AOAC International*, 80: 1215-1219.
- Charoenpornsook, K. and P. Kavisarasai (2006). Mycotoxins in animal feedstuffs of Thailand. *KMITL Sci. Tech. J.*, 6: 25-28.
- Davis, N. D., J. W. Dickens, R. L. Freie, P. B. Hamilton, O. L. Shotwell and T. D. Wylie (1980). Protocols for surveys, sampling, post-collection handling and analysis of grain samples involved in mycotoxin problems. *Journal of AOAC*. 63: 5-103.
- Dersjant-Li, Y., M. W. A. Verstegen and W. J. J. Gerrits (2003). The impact of low concentrations of aflatoxin, deoxynivalenol or fumonisin in diets on growing pigs and poultry. *Nutrition Research Reviews*, 16: 223-239.
- Diaz, G. J., E. Calabrese and R. Blain (2008). Aflatoxicosis in Chickens (*Gallus gallus*): An example of Hormesis? *Poultry Science*, 87: 727-732.
- Diaz, G. J., N. S. Perilla and Y. Rojas (2001). Occurrence of aflatoxins in selected Colombian foods. *Mycotoxin Research*, 17: 5-20.
- Ghahri, H., R. Habibian and M.A. Fam (2010). Effect of sodium bentonite, mannan oligosaccharide and humate on performance and serum biochemical parameters during aflatoxicosis in broiler chickens. *Global Veterinaria*, 5: 129-134.
- Hanif, N. Q., M. Naseem, S. khaton and N. Malik (2006). Prevalence of mycotoxins in poultry rations. *Pakistan Journal of Science and Industrial Research*, 49: 120-124.
- Hanif, N. Q., G. Muhammad, M. Siddique, A. Khanum, T. Ahmad, J. A. Gadahai and G. Kaukab (2008). Clinico-pathomorphological, serum biochemical and histological studies in broilers fed ochratoxin A and a toxin deactivator (Mycifix® PlusMTV INSIDE). *British Poultry Science*, 49: 632-642.
- Hussain Z., M. Z. Khan, A. Khan, I. Javid, M. K. Salem, S. Mahmood and M. R. Asi (2010). Residues of aflatoxin B<sub>1</sub> in broiler meat: Effect of age and dietary aflatoxin B<sub>1</sub> levels. *Food and Chemical Toxicology*, 48: 3304-3307.
- Khan, W. A., M. Z. Khan, A. Khan and I. Hussain (2010). Pathological effects of aflatoxin and their amelioration by vitamin E in white leghorn layers. *Pak. Vet. J.*, 30: 155-162.
- Ibrahim, I. K., A. M. Shareef and K. M. T. Al-Joubory (2000). Ameliorative effects of sodium bentonite on phagocytosis and Newcastle disease antibody formation in broiler chickens during aflatoxicosis. *Research in Veterinary Science*, 69: 119-2.
- Krnjaja, V., L.J. Stojanovic, R. Cmiljanic, S. Trenkovski and D. Tomasevic (2008). The presence of potentially toxigenic fungi in poultry feed. *Biotechnology in Animal Husbandry*, 24: 87-93
- Maqbool, U., M. Ahmad, Anwar-ul-Haq and M. M. Iqbal (2004). Determination of aflatoxin B<sub>1</sub> in poultry feed and its components employing enzyme-linked immunosorbent assay (ELISA). *Toxicol. and Environ. Chem.*, 86: 213-218.
- Mngadi, P.T., R. Govinden and B. Odhav (2008). Co-occurring mycotoxins in animal feeds. *African Journal of biotechnology*, 7: 2239-2243.
- Moalem, S.H.H., G. Azizi and M. Azarmi (2010). Prevalence of aflatoxin B<sub>1</sub> in feedstuffs in northern Iran. *Global Veterinaria*, 4: 144-148.
- Moss, M. O. (2002). Mycotoxin review -1. *Aspergillus and penicillium*. *Mycologist*, 16: 116-119.

- Oguz, H., H. H. Hadimli, V. Kurtoglu and O. Erganis (2003). Evaluation of humoral immunity of broilers during chronic aflatoxin (50 and 100 ppb) and clinoptilolite exposure. *Revue Med. Vet.*, 154: 483-486.
- Ortatatli, M., H. Ogus, F. Hatipoglu and M Karaman (2005). Evaluation of pathological changes in broilers during chronic aflatoxin (50 and 100 ppb) and clinoptilolite exposure. *Research in veterinary Science*, 78: 61-68.
- Qazi, J. I. and Z. Fayyaz (2006). Aflatoxin contaminated foods and health risk perspective for Pakistani population. *Mycopath.*, 4: 27-34.
- Richard, J. (2000). Sampling and sample preparation for mycotoxin analysis. *Romer lab's guide to mycotoxins vol. 2*. Romer™ Labs, Inc. 1301 Stylemaster Drive, Union, MO63084-1156. : 15.
- Shi, Y., Z. Xu, Y. Sun, C. Wang and J. Feng (2009). Effects of two different types of montmorillonite on growth performance and serum profiles of broiler chicks during aflatoxicosis. *Turk. J. Vet. Anim. Sci.*, 33: 15-20.
- Shareef, A.M. (2010). Molds and mycotoxins in poultry feeds from farms of potential mycotoxicosis. *Iraqi Journal of Vet. Sci.*, 24: 17-25.
- Shareef, A. M., O. S. Bayon, and T. S. Qubih (2008). Correlation between broiler aflatoxicosis and European production efficiency factor. *Iraqi Journal of Vet. Sci.*, 22: 49-55.
- Sultana, N. and N.Q. Hanif (2009). Mycotoxin contamination in cattle feed and feed ingredients. *Pak. Vet. J.*, 29:211-213.
- Thompson, C. and S. E. Henke (2000). Effect of climate and type of storage container on aflatoxin production on corn and its associated risks to wildlife species. *Journal of Wildlife Diseases*, 36: 172-179.
- Yunus, A. W., M. K. Nasir, T. Aziz and J. Bohm (2009). Prevalence of poultry diseases in district Chakwal and their interaction with mycotoxicosis: 2. Effects of season and feed. *The Journal of Animal and Plant Sciences*, 19: 1-5.