

MOLECULAR CHARACTERIZATION AND PHYLOGENETIC ANALYSIS OF *MYCOPLASMA SYNOVIAE* ISOLATED FROM CHICKEN

A. Rasool¹, A. A. Anjum¹, M. Rabbani², M. Lateef⁴, F. Akhter², H. Afroz³, J. Muhammad² and M. Nawaz¹

¹Department of Microbiology, University of Veterinary and Animal Sciences, Lahore, Pakistan.

²University Diagnostic Laboratory, University of Veterinary and Animal Sciences, Lahore, Pakistan.

³Veterinary Research Institute, Zarar Shaheed Road, Lahore, Pakistan.

⁴Department of Parasitology, University of Veterinary and Animal Sciences, Lahore, Pakistan.

Corresponding E-mail: aftab.anjum@uvas.edu.pk

ABSTRACT

Mycoplasma synoviae is one of the paramount causes of economic losses for poultry industry. Present study was carried out with the aim to isolate, identify and characterize *M. synoviae* from suspected chicken. Tracheal and cloacal swabs samples (n=500), collected from suspected chicken, were cultured on Modified Frey's medium for isolation of *Mycoplasma*. Preliminary identification of *Mycoplasma* was done by culturing and biochemical characterization. Partial 16S rRNA was amplified and sequenced for confirmation and molecular identification of *M. synoviae*. For strain identification and phylogenetic analysis, partial amplification and sequence of *vlhA* gene was performed. Out of 500 samples, *Mycoplasma* was successfully isolated from 40 samples. Out of 40 mycoplasma isolates, 20 were identified as *M. synoviae* by specie specific 16S ribosomal RNA gene, partial amplification (~213 bp). The *vlhA* gene was also successfully partially amplified (~375 bp) and sequenced from five isolates. GenBank accession numbers of reported local isolates in this study are KJ130523-KJ130542 and KP316017-KP316021 respectively.

Key words: Mycoplasma, fried egg colonies, 16S rRNA, *vlhA* gene

INTRODUCTION

Mycoplasma is one of the major respiratory tract pathogen of chicken, causing considerable losses to poultry industry (Dufour-Gesbert *et al.*, 2006). *Mycoplasma synoviae* and *Mycoplasma gallisepticum* are considered as most pathogenic avian strains (Yilmaz *et al.* (2011). *M. synoviae* mostly infects upper respiratory tract as a subclinical infection but if it combines with *M. gallisepticum* it may cause air sac lesions (Silva *et al.*, 2008). *M. synoviae* causes heavy losses due to reduction in egg production and growth rate, loss of egg production, egg shell abnormalities, low hatchability and lame birds culling (Vogel *et al.*, 2008; Peebles *et al.*, 2011). *M. synoviae* transmission occurs both vertically and horizontally. *M. synoviae* is transferred through infectious aerosols which are generated by coughing and sneezing of infected birds, contaminated feed, water, feather, personal contact and direct contact of susceptible birds with infected chickens (Marois *et al.*, 2005; Butcher and Jacob, 2009). Immune suppression in birds enhances the spread of infection (Dhondt *et al.* (2007). True and correct diagnosis is required to prevent infection dissemination (Seifi and Shirzad, 2012).

For the screening of *M. synoviae* in chicken flocks rapid slide agglutination test (RSA) and Enzyme linked immunosorbent assay (ELISA) are widely used. Species-specific 16S ribosomal RNA gene based PCR is also used to detect the infection but due to conserved

nature of gene it does not allow strain differentiation of *M. synoviae*. Although there is only one serotype of *M. synoviae* present but heterogeneity among *M. synoviae* strains is observed (Kleven, 2003) which can be determined by amplification and sequencing the *vlhA* gene. *M. synoviae* *vlhA* gene is expressed as two products including hemagglutinin and prolipoprotein (Noormohammadi *et al.*, 1998). The hemagglutinin has high frequency of antigenic and phase variation (Noormohammadi *et al.*, 2000; Kleven, 2003). Recombination of *vlhA* gene at 3' end with one of the pseudo genes can produce antigenic variants which may lead to evasion of host immune system. Although *vlhA* gene 5' end is relatively conserved and present as a single copy, but pseudo genes can replace its downstream region (Noormohammadi *et al.*, 2000). For the isolation and control of prevalent *M. synoviae* it is very important to characterize its isolates up to its strain level for better understanding. For PCR based strain identification the upstream region of 5' of *vlhA* gene is used in outbreak and epidemiological studies (Hong *et al.*, 2004; Hammond *et al.*, 2009).

Although there are many studies throughout the world which report the isolation and molecular characterization of *M. synoviae* from chicken, only a few reports are available from Pakistan. Therefore, present study was designed for isolation, and identification of *M. synoviae* from dubious chicken, partial amplification of *vlhA* and 16S rRNA gene and sequencing of amplicons

for phylogenetic analysis to determine the heterogeneity of *M. synoviae* isolates.

MATERIALS AND METHODS

A total of 500 samples including tracheal/cloacal swabs (n=250) and morbid materials such as joints, trachea and lungs (n=250) were collected from chicken suspected with *Mycoplasma* infection. Trachea and cloacal cleft swabs were transported in Aimes transport media to University Diagnostic Laboratory (UDL), University of Veterinary and Animal Sciences (UVAS), Lahore, Pakistan. Morbid materials including trachea, air sacs, lungs and joints were collected and transported to lab in polythene bags with proper labels and case history.

Isolation of Mycoplasma: For the isolation of Mycoplasma, all samples were cultured in Frey's media (Lavric *et al.*, 2007). Each set of five samples were cultured as a unit. Tracheal/cloacal swabs were cultured directly. Joints were opened, swabbed from internal surface of joint cavity and cultured. Other organs such as trachea and lungs were grinded with phosphate buffer saline, centrifuged at 1000 rpm for 20 minutes and supernatants were filtered through a syringe filter (0.22µm) and cultured at 37 °C in 5% carbon dioxide. Cultures were observed for any change in color and turbidity. Initial cultures were filtered with or without showing any growth through 0.22µm filter and sub cultured in modified Frey's broth and agar. Cultures were observed daily for 21 days for any change in color, turbidity and presence of typical fried egg shaped colonies on agar plates, to be declared as negative.

Biochemical identification: Culture was identified as mycoplasma by biochemical characterization and microscopic examination. For microscopic examination the impression smears were prepared from fried egg colonies, stained by Giemsa stain and observed at 1000X magnification using bright field microscope. The biochemical tests performed for suspected cultures were glucose fermentation, arginine hydrolysis and tetrazolium dye reduction test (Ehtisham *et al.*, 2011).

Molecular characterization: The isolates, identified as Mycoplasma on the basis of morphological, cultural and biochemical characteristics, were confirmed as *M. synoviae* by specie specific 16S r DNA PCR. Briefly, DNAs of all isolates were extracted by QAIKEN DNA extraction kit. Partial 16S DNA was amplified by using *M. synoviae* specific primer pair MS-F; 5'-GAG-AAG-CAA-AAT-AGT-GAT-ATC-A-3' and MS-R: 5'CAG-TCG-TCT-CCG-AAG-TTA-ACA-A-3' (Lauerman, 1998). Reaction mixture (25 µL) had ~20 n mol of genomic DNA and 25 p moles of reverse and forward primer. In Amplification cycle Initial denaturation occurs at 95 °C for 5 minutes, denaturation at 95 °C, 35 cycles of

30 second each, annealing at 49°C, one minute extension at 72 °C, and final extension at 72°C for 5 minutes was done. Amplicons (213 bp) were resolved on agarose gel (2 %), stained with ethidium bromide (0.05 µg/ml) and visualized in UV illuminator and gel documentation system (Bio-Rad, USA)

Partial amplification of N-terminal of vlhA gene: MS vlha- F; 5'-GGC-CAT-TGC-TCC-TAC-TGT-TAT- 3' MS vlha- R; 5'-AGT-AAC-CGA-TCC-GCT-TAA-TGC-3' following Wetzel *et al.*, (2010) were used to amplify vlhA gene N-terminal using 56° C annealing temperatures. Amplicons (375bp) were resolved on agarose gel (2%), stained with ethidium bromide (0.05 µg/ml) and visualized in UV illuminator and gel documentation system (Bio-Rad, USA)

Sequencing and Phylogenetic analysis: Partial 16S r DNA amplicons (213 bp) and vlhA amplicons (385 bp) were sequenced by Sanger method. Sequence results were analyzed using Bio Edit software and sequence homologies were determined by Clustal W. Gen Bank accession numbers for all the sequences were obtained from NCBI. Phylogenetic trees were constructed by MEGA 6 version, using NJ method with 1000 bootstrap value.

RESULTS

Total of 500 samples were cultured as 100 units (each unit contained five samples). Out of 100 units, growth of *Mycoplasma* was observed from 40 units (40%) including 20 units of tracheal /cloacal swabs, 15 units of joints and 05 units of samples of trachea, lungs and air sacs. Growth of *Mycoplasma* was observed as typical egg fried shaped colonies on Frey's agar (Fig. 1a). Impression smears prepared from egg fried colonies were stained with Giemsa stain and *Mycoplasma* specific pleomorphic coccoid bodies were observed under microscope at 1000X magnification (Fig. 1b). All isolates (n=40, MYC01-MYC40) were positive for glucose fermentation while negative for arginine hydrolysis. Isolates showed variable results for tetrazolium reduction. Out of 40 Mycoplasma isolates, 20 isolates showed *M. synoviae* specific amplification (amplicon size ~213 bp) of 16S rDNA by PCR (Fig 2). Analysis of partial sequences also revealed that all these 20 isolates were *M. synoviae*. GenBank Accession numbers of these isolates are KJ130523-KJ130542. Phylogenetic tree of *M. synoviae* isolates (n=20), on partial 16S r DNA sequences, revealed that all isolates had close genetic relatedness (> 99% similarity (Fig. 3). Furthermore, phylogenetic tree based on 16S rDNA (Partial amplification) also revealed that *M. synoviae* isolated in this study had close relatedness with isolates from Iran, India and Egypt. The vlhA gene was successfully amplified (figure 04) from 05 isolates. Sequences of vlhA

gene (Partial amplification) were submitted to NCBI. GenBank accession numbers are KP316017-KP316021. Phylogenetic tree, constructed on the basis of sequences of *vlhA* gene amplicons (partial), revealed that isolates had similarities with isolates from Europe and USA (Fig. 5).

DISCUSSION

Mycoplasma synoviae is considered as one of the important leading agent to be responsible for economic losses in poultry sector. Poultry flocks are generally screened for *M. synoviae* by rapid slide agglutination test, enzyme linked immunosorbent assay and polymerase chain reaction throughout the world (Hess *et al.*, 2007). Isolation of *M. synoviae* is considered as touchstone for confirmation of infection. Isolation of *M. synoviae* is a prerequisite for its molecular characterization, phylogenetic analysis and possible source tracking (Bayatzadch *et al.*, 2011; Buim *et al.*, 2009). Isolation of *M. synoviae* is not preferred for screening of large flocks because of difficulties in its culture and low recovery rate. Phenotypic and genotypic characterization of *M. synoviae* helps in future diagnostic and epidemiological studies (Moreira *et al.*, 2015).

M. synoviae being a fastidious organism tends to disappear from the lesions after few weeks so its isolation and recovery rate is dependent on the time and site of sampling. In present study different types of samples including tracheal/cloacal cleft swab, trachea, air sacs, lungs and joint were collected to enhance the recovery rate of *M. synoviae* (Hennigan *et al.*, 2012; Ehtisham *et al.*, 2011; Pourbakhsh *et al.*, 2010). *M. synoviae* were successfully isolated on Frey's media (pH 7.8) at 37 °C in 10% CO₂ and high humidity. Similar growth conditions have been used previously by Tebyanian *et al.*, 2014). Cultured samples, with or without any indication of growth, were sub cultured in fresh Frey's medium after filtering through 0.22 µM syringe filters. Previous studies have used filtration of sample or initial culture through filter paper of 0.45µM pore size to remove the contaminants (Bayatzadch *et al.*, 2011; Tebyanian *et al.*, 2014). In present study, filtration was done through a filter paper of 0.2 µM pore size, which proved to be more efficient for the isolation and purification of *M. synoviae*. Growth of *mycoplasma* was observed by change of phenol indicator in Frey's media from red to orange and yellow as mycoplasma produces acid metabolites and lowers the pH of media. Low pH is injurious for mycoplasma survival, so it is very important to remove the positive samples from incubation and kept under refrigeration for further processing. Overall recovery rate of Mycoplasma (40%), reported in this study, is in meeting of minds

with previous studies (Nikfarjam and Farzaneh, 2012; Lavric *et al.*, 2007; Bencina *et al.*, 2001). *Mycoplasma* showed fried egg shaped colonies on solid media (Ehtisham *et al.*, 2011; Kleven and Ferguson-Noel, 2008; Kleven, 2003).

Although 16S rDNA is the most conserved region in bacterial genome, it still contains specie specific signature sequences which is used as an momentous specie identification tool (Bayatzadeh *et al.*, 2013; Lobova *et al.*, 2012; Perez *et al.*, 2011; Ehtisham *et al.*, 2011; Jarquin *et al.*, 2009). *M. synoviae* specific amplification revealed that out of 40 isolates of Mycoplasma, 20 (50%) was *M. synoviae*. Sequencing of specific PCR amplicons (213 bp) was used for confirmation and phylogenetic analysis. All *M. synoviae* isolates showed more than 99 % similarity to each other and other *M. synoviae*. Neighbor-Joining method is used to extrapolate evolutionary history (Saitou and Nei, 1987). Optimal tree with branch length sum of 0.54344747 is shown. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to deduce the phylogenetic tree. The evolutionary distances were reckoned by the Maximum Composite Likelihood method and were in units of number of base substitutions per site. Variation rate among sites was modeled with a gamma distribution (shape parameter = 3). Phylogenetic analysis on 16S rDNA partial amplicons sequences includes a total of 5 sequences reported in this study and 07 outgroup. All positions with missing data and gaps were precluded. There were total of 197 positions in the final dataset.

M. synoviae partial amplification of *vlhA* gene is useful for *M. synoviae* strain identification (Dijkman *et al.*, 2014). Out of 20 isolates of *M. synoviae*, *vlhA* gene was successfully partially amplified from 05 (20%) isolates. These results are in agreement with Ley, (2008) and Hammond *et al.*, (2009). For *vlhA* partial amplification primers used by Wetzel *et al.*, (2010) were used. Phylogenetic tree, based on *vlhA* gene partial sequences revealed the evolutionary history of the isolates reported in this study. The Maximum Composite Likelihood method is used to reckon the evolutionary distances and the number of base substitutions per site is used as unit. The analysis involved 16 nucleotide sequences with Codon positions 1st+2nd+3rd+Noncoding. All positions with missing data and gaps were closed out. There were total of 298 positions in the final data set. MEGA 6 is used to analyze evolution (Tamura *et al.*, 2013).

M. synoviae culture and its isolation is not an easy job but it is touchstone in the mycoplasma infection confirmation. Its phenotypical and genotypical characterization can help in future diagnostic and epidemiological studies.




Figure 1: (A) Fried egg colonies of Mycoplasma, (B) Giemsa stained colonies of Mycoplasma




Figure 2: Partially amplified 16S ribosomal RNA gene of *M. synoviae*




Figure 3: Partially amplified vlnA gene of *M. synoviae*




Figure 4. Phylogenetic tree of *M. synoviae* isolates based on 16S ribosomal RNA gene, partial sequence.




Figure 5. Phylogenetic tree of *M. synoviae* isolates based on vlh A gene, partial sequence

REFERENCES

Bencina, D., M. Dorbnic Valic, S. Horvar, M. Narat, S.H. Kleven, and P. Dovc (2001). Molecular basis of the length variation in the terminal part of *Mycoplasma synoviae* haemagglutinin. FEMS. Microbial. LETT. 203: 115-123.

Bayatzadeh, M.A., S.A. Pourbakhsh, A.R. Homayounimehr, A. Ashtari and A.R. Abtin (2011). Application of culture and polymerase

chain reaction (PCR) methods for isolation and identification of *Mycoplasma synoviae* on broiler chicken farms. Arch. Razi Inst. 66(2): 87-94.

Bayatzadeh, M.A., S.A. Pourbakhsh, A.R. Homayounimehr, A. Ashtari and A.R. Abtin (2013). Phylogenetic analysis of *Mycoplasma synoviae* isolated from commercial Iranian chicken farms compared with other GeneBank isolate sequences based on 16S rRNA gene. Avian Biol. Res. 6(3): 233-238.

- Buim, M.R., E. Mettifogo, J. Timenetsky, S. Kleven and A.J.P. Ferreira (2009). Epidemiological survey on *Mycoplasma gallisepticum* and *M. synoviae* by multiplex PCR in commercial poultry. *Pesqui. Vet. Brasil.* 29: 552-556.
- Butcher, G.D. and J.P. Jacob (2009). Respiratory diseases: common poultry diseases. *Poult. Sci.* 47: 6-7.
- Dhondt, A.A., K.V. dhond, D.M. Hawley, and S. Jennelle (2007). Mission of *Mycoplasma gallisepticum* in house finches by fomites. *Avian Pathol.* 36: 205-208.
- Dijkman, R., A. Feberwee and W.J. Landman (2014). Variable lipoprotein haemagglutinin (vlhA) gene sequence typing of mainly Dutch *Mycoplasma synoviae* isolates: comparison with vlhA sequences from Genbank and with amplified fragment length polymorphism analysis. *Avian Pathol.* 43: 465-472.
- Dufour-Gesbert, F.A., C. Dheilly, I. Marois, and I. Kempf (2006). Epidemiological study on *Mycoplasma synoviae* infection in layers. *Vet. Microbiol.* 114: 148-154.
- Ehtisham-ul-haq, S., S.U. Rehman, M. Siddique and A.S. Qureshi (2011). Involvement of *Mycoplasma synoviae* in respiratory distress cases of broiler. *Pakistan Vet. J.* 31(2): 117-119.
- Hammond, P.P., A.S. Ramirez, C.J. Morrww, and J.M. Bardbury (2009). Development and evaluation of an improved diagnostic PCR for *Mycoplasma synoviae* using primers located in the haemagglutinin encoding gene vlhA and its value for strain typing. *Vet. Microbiol.* 136(1-2): 61-68.
- Hennigan, S.L., J.D. Driskell, N.F. Noel, R.A. Dluhy, Y. Zhao, R.A. Tripp and D.C. Krausea (2012). Detection and differentiation of avian mycoplasmas by surface-enhanced raman spectroscopy based on a silver nanorod array. *Appl. Envir. Microbiol.* 78: 1930-1935.
- Hess, M., C. Neubauer and R. Hackl (2007). Interlaboratory comparison of ability to detect nucleic acid of *Mycoplasma gallisepticum* and *Mycoplasma synoviae* by polymerase chain reaction. *Avian Pathol.* 36:127-133.
- Hong, Y., M. Garacia, V. Leiting, D. Bencina, L. Dufour-Zavala, G. Zavala, and S.H. Kleven (2004). Specific detection and typing of *Mycoplasma synoviae* strain in poultry with PCR and DNA sequencing analysis targeting the haemagglutinin encoding gene vlhA. *Avian Dis.* 48: 606-616.
- Jarquin, R., J. Schultz, I. Hanning and S.C. Ricke (2009). Development of a real-time polymerase chain reaction assay for the simultaneous detection of *Mycoplasma gallisepticum* and *Mycoplasma synoviae* under industry conditions. *Avian Dis.* 53: 73-77.
- Kleven, S.H. (2003). *Mycoplasma synoviae* Infection. In: Diseases of poultry, 11th ed. Y M Saif (Ed.). Iowa State University Press, Ames, IA. Pp: 756–766.
- Kleven, S.H. and N. Ferguson-Noel (2008). *Mycoplasma synoviae* infection. In: Saif YM, editor. Diseases of Poultry. Ames, Iowa: Blackwell Publishing Ltd. Pp: 845-856.
- Lauerma, L.H. (1998). *Mycoplasma* PCR assays. In: nucleic acid amplification assays for diagnosis of animal diseases. Am. Assoc. Vet. Lab. Diag. Pp: 41-42.
- Lavric, M., D. Bencina, S. Kothlow, B. Kasper, and M. Narat (2007). *Mycoplasma synoviae* lipoprotein MSPB, the N-terminal part of VlhA haemagglutinin, induces secretion of nitric oxide, IL-6, IL-1 beta in chicken macrophages. *Vet. Microbiol.* 121: 278-287.
- Ley, D.H. (2008). *Mycoplasma gallisepticum* Infection. In: Saif YM, editor. Diseases of Poultry. 12. Ames: Blackwell Publishing Ltd. Pp: 807-834.
- Lobova, D., L. Kohoutova, D. Molinkova, K. Rosenbergova, O. Kubicek, and V. Celer (2012). Prevalence of etiological agents of selected respiratory infections in chicken and turkey farms in the Czech Republic. *Vet. Med.* 57: 125-132.
- Marois, C., J. Picault, M. Kobisch, and I. Kempf (2005). Experimental evidence of indirect transmission of *Mycoplasma synoviae*. *Vet. Res.* 36: 759-769.
- Moreira, R., F. Chenlo, S. Arufe and S.N. Rubinos (2015). Physicochemical characterization of white, yellow and purple maize flours and rheological characterization of their doughs. *J. Food Sci. Technol.* 52: 7954-7963.
- Nikfarjam, L. and P. Farzaneh (2012). Prevention and detection of *Mycoplasma* contamination in cell culture. *Cell J.* 13: 203-212.
- Noormohammadi, A.H., P.F. Markham, M.F. Duffy, K.G. Whithear, and G.F. Browning (1998). Multigene families encoding the major hemagglutinins in phylogenetically distinct mycoplasmas. *Infect. Immun.* 66: 3470–3475.
- Noormohammadi A.H., P.F. Markham, A. Kanci, K.G. Whithear, and G.F. Browning (2000). A novel mechanism for control of antigenic variation in the haemagglutinin gene family of *Mycoplasma synoviae*. *Mol. Microbiol.* 35: 911–923.
- Peebles, E.D., S.W. Park, S.L. Branton, P.D. Gerard, and S.K. Womack (2011). Dietary poultry fat, phytase, and 25-hydroxycholecalciferol influence the digestive and reproductive organ characteristics of commercial layers inoculated before or at the onset of lay with F-strain

- Mycoplasma gallisepticum* 1, 2. Poul. Sci. 90 (4): 797-803.
- Pérez, C., K. Hausteijn, Z. Janjic, O. Jorba, N. Huneus, J.M. Baldasano, T. Black, S. Basart, S. Nickovic, R.L. Miller, J.P. Perlwitz, M. Schulz, and M. Thomson (2011). Atmospheric dust modeling from meso to global scales with the online NMMB/BSC-Dust model — Part 1: Model description, annual simulations and evaluation. *Atmos. Chem. Phys.* 11: 13001-13027.
- Pourbakhsh, S.A., G.R. Shokri, M. Banami, F. Elhamnia and A. Ashtari (2010). Detection of *Mycoplasma synoviae* in broiler breeder farms of Tehran province using PCR and culture methods. *Arch. Razi Institute.* 65(2): 75-81.
- Saitou, N. and M. Nei (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4: 406-425.
- Seifi, A. and M.R. Shirzad (2012). Incidence and risk factors of *Mycoplasma synoviae* infection in broiler breeder farms of Iran. *Vet. World.* 5(5): 265-268.
- Silva, R.C.F., E.R.D. Nascimento, V.L.A. Pereira, M.L. Barreto and M.G.D. Nascimento (2008). *Mycoplasma synoviae* infection on Newcastle disease vaccination of chickens. *Braz. J. Microbiol.* 39: 384-389.
- Tamura, K., G. Stecher, D. Peterson, A. Filipski and S. Kumar (2013). MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol. Biol. Evol.* 30: 2725-2729.
- Tebyanian, H., S.H. Mirhosseiny, B. Kheirkhah, M. Hassanshahian and H. Farhadian (2014). Isolation and Identification of *Mycoplasma synoviae* From Suspected Ostriches by Polymerase Chain Reaction, in Kerman Province, Iran. *Jundishapur J. Microb.* 7: e19262.
- Vogel, G.A., S. plaickner, L. Szathmary, R. Stipkovitis, Rosengarton and M.P. Szostak (2008). *Mycoplasma gallisepticum* invades chicken erythrocytes during infection. *Infect. Immun.* 76: 71-77.
- Wetzel, A.N., M. Kristina, Lefevre and Z. Raviv (2010). Revised *Mycoplasma synoviae* V1ha PCRs. *Avian Dis.* 54: 1292-1297.
- Yilmaz, F., A. Timurkaan, A. Kilic, H. Kalender and U. Kilinc (2011). Detection of *Mycoplasma synoviae* and *Mycoplasma gallisepticum* in chicken by immune histochemical, PCR and culture methods. *Rev. Med. Vet.* 162(2): 79-86.