

EVALUATION OF ANTIOXIDANT AND ANTIHYPERLIPIDEMIC ACTIVITY OF INDIAN GOOSEBERRY (*Emblica officinalis*) FRUIT IN HIGH FAT-FED RABBITS

S. Zahid¹, K. M. Anjum^{2*}, M. S. Mughal³, A. Yaqub³ and M. Yameen⁴

¹Discipline of Zoology, ²Department of Wildlife & Ecology, Faculty of Fisheries & Wildlife, University of Veterinary & Animal Sciences, Lahore-Pakistan; ³Department of Zoology, Government College University, Lahore-Pakistan

⁴Department of Applied Chemistry & Biochemistry, Government College University, Faisalabad-Pakistan

*Corresponding Author: Khalid.mahmood@uvas.edu.pk

ABSTRACT

Hyperlipidaemia leads to various diseases including cardiovascular and diabetes mellitus, etc. Among the many reasons for the abnormal lipid profile, high fat intake and sedentary life style are the main reasons for hypercholesterolemia. In the present study, a total of 25 ($n=25$) rabbits were used and were randomly divided into five groups which received different treatments. Group I served as control, group II received a measured quantity of *Emblica officinalis* (EO) powder, group III received high fat diet (HFD), group IV received HFD and EO powder and group V received HFD and the standard drug atorvastatin for eight weeks. To investigate the effects of EO powder, blood and tissue (liver and kidney) samples were taken at the end of the eight weeks of experimentation and serum was evaluated for various blood plasma lipid parameters and antioxidant enzymes of liver and kidney. There was a significant increase in all the lipid parameters except for HDL in the group of rabbits receiving HFD; HDL level decreased significantly ($P < 0.05$). High fat intake may have resulted in the production of free radicals which created an oxidative stress and lipid peroxidation ultimately leading to abnormal lipid profile. Antioxidant enzymes, such as Superoxide dismutase (SOD), Catalase (CAT) and Glutathione (GSH) were significantly reduced in the HFD group indicating clearly that high fat diet leads to oxidative stress. *Emblica officinalis* relieved the oxidative stress of liver and kidney tissues in group IV indicating that EO powder has the potential to inhibit the oxidative stress and lipid peroxidation because of poly-phenols, flavonoids, ascorbic acid and other chemicals present in it. Further, it is concluded that although statins are useful in reducing blood plasma lipid profile, they simultaneously have many side effects. The findings of the present study reveal that EO powder is useful in reducing hyperlipidaemia and tissue damage.

Keywords: *Emblica officinalis*, Atorvastatin, Hyperlipidemia, Lipid Profile, Medicinal Plants.

INTRODUCTION

High blood cholesterol level is the major reason for various diseases including nervous and cardiovascular disorders (Goldstein and Brown, 2015). There are a number of reasons for this hyperlipidemic condition including high fat diet, sedentary lifestyle, genetics, kidney problems and liver disease. Several factors, such as increased age, lack of exercise, obesity, smoking, excessive alcohol intake, diabetes, low thyroid and Cushing's syndrome increase the risk to be hyperlipidemic (Kawasumi *et al.*, 2014). There are several epidemiological findings that indicate high blood cholesterol, clotting factors and oxidation of lipoproteins lead to the formation and rupture of atherosclerotic plaque (Cappelli-Bigazzi *et al.*, 1997), decreasing the oxygenated blood flow towards vital organs leading to angina, heart attack, stroke and other serious complications (Herron *et al.*, 2004; Varbo *et al.*, 2014).

Cholesterol lowering drugs of different classes (Statins, Niacin, Bile-acid binding resins, Omega-3 fatty acids, Fibrates, Cholesterol absorption inhibitor, etc.) are

frequently used to avoid damages of hyperlipidaemia. Mostly these medicines reduce low density lipoproteins (LDL) level and some also increase the high density lipoproteins (HDL) level in blood circulation through different mechanisms (Illingworth, 1987). Although these drugs are very beneficial to lower the bad lipid profile parameters, they also have adverse side effects including memory loss, depression, insomnia, itching and headache, etc. (Kureishi *et al.*, 2000; Pahan, 2006).

Man has been using the natural herbal drugs since centuries. Researchers are trying their best to investigate natural plants for their role for the cure of different ailments (Anjum *et al.*, 2014). There are some evidences that some plants have immense potential for medicinal use. The use of several plants has been reported for the treatment of hyperlipidaemia. Natural products must preferably be used to lower the blood lipids. These products not only rescue the body from hyperlipidaemia but also improve the function of other body organs. For example, the fruit of *Emblica officinalis* (EO) is used to manage the blood glucose levels but simultaneously it improves the digestive system and

protects the body from harmful microbes (Ahmad *et al.*, 1998; Khan, 2009).

Emblica officinalis (EO), a deciduous ornamental tall tree (approximately 60 ft. tall), is a member of family Euphorbiaceae, locally known as “amla”. It is a traditional medicinal plant native to tropical parts of Southeast Asia. Due to its medicinal and economic importance, it is cultivated commercially in India, China and some other countries (Sairam *et al.*, 2002). All parts of this plant are reported to be used for preparation of various herbal medicines to cure various ailments. Decoction and infusion of leaves and seeds of this plant are commonly used (Kumar *et al.*, 2012b). These extracts and distillates of leaves and seeds are reported to have bactericidal, anti-inflammatory and antipyretic properties and are used as natural mouth wash and also applied against cold, dysentery, fever and urine retention (Jain *et al.*, 2014). The flowers and roots are used as refrigerant, aperient and emetic (Abbasi *et al.*, 2010) and their extracts are used for the treatment of gonorrhoea, diarrhoea and mouth inflammations (Kumar *et al.*, 2012a). Seeds and seed oil of this plant are used to cure asthma, skin complications and bronchitis (Jain *et al.*, 2015; Khan, 2009).

Emblica officinalis fruit is considered diuretic and laxative while fruit juice is used to cure jaundice, dyspepsia and coughs (Kumar *et al.*, 2012b). The powdery paste of the fruit is an expectorant and also used for the treatment of various diseases like indigestion, heart problems and nasal congestion, etc. (Jain *et al.*, 2014). The juice from leaves and fruit act as antimicrobial agent to cure sores and wounds (Ahmad *et al.*, 1998). Leaves and fruit of EO plant are reported to be rich in tannin, ascorbic acid, malic acid, flavonoids and ellagic acid, etc. (Scartezzini *et al.*, 2006). There are several reports that various medicinal plants possess antioxidative properties. Antioxidant enzymes protect body and tissues by various mechanisms. Superoxide dismutase (SOD) converts harmful “superoxide anions (O₂⁻)” to H₂O₂ and hydrogen peroxide is catalysed by catalase (CAT) to water and O₂. Further, glutathione detoxify H₂O₂ and other organic hydro-oxides, thus protect tissues and relieves the oxidative stress (Bazmandegan *et al.*, 2017). Flavonoids are widely distributed in the plant phenolic compounds, known to have more than 4000 kinds of individual substances (Qasim *et al.*, 2017). Plants use these flavonoids to prevent oxidative damage by suppressing the free radicals and reactive oxygen species (ROS) produced by sunlight. Dietary polyphenols enhance the stability of LDL against oxidation (Scartezzini *et al.*, 2006). Several studies have reported that oxidized LDL, in turn, plays an important role in atherosclerosis and cardiovascular disease (CVD). Epidemiological studies have proven that intake of antioxidant rich foods are inversely proportional to the CVD (Shyamala *et al.*, 2003). Most of the phenolic

compounds have potential to act as antioxidants because they possess conjugated ring structure and the hydroxyl group. Free radicals, hydrogen peroxide leads to oxidative stress which is cytotoxic to the tissues and antioxidants from flavonoids protect tissues and make complexes and hydrogenate free radicals and convert them to neutral compounds (Binti Anzian *et al.*, 2017).

Keeping in view the broad beneficial aspects of EO fruit, the present study was conducted to evaluate the antioxidative and hypolipidemic potential of this fruit in high fat fed rabbits. Furthermore, anti-hyperlipidemic and antioxidative efficiency of EO fruit powder was also compared with a standard cholesterol lowering drug.

MATERIALS AND METHODS

Animals: The study was conducted at the Department of Wildlife & Ecology, University of Veterinary and Animal Sciences (UVAS), Lahore, Pakistan. Twenty five large rabbits (*Oryctolagus cuniculus*) were used for the study. Their mean ages was 50±10 days and mean body weight was 1.75±0.25 Kg either sex. Animals were acclimatized for 10 days in the University animal enclosures under standard conditions before the start of the experiment. The animals were fed on normal diet consisting of fodder and grams in sufficient quantity throughout the duration of the experiment and water was provided *ad libitum*.

***Emblica officinalis* (EO):** EO fruit was purchased from local market, identified and authenticated by a professional botanist, washed with water, air dried and ground along with the seeds.

Atorvastatin: Atorvastatin is a standardized medication for the reduction of circulatory blood cholesterol level. This drug was purchased from a local medical store. Atorvastatin belongs to the broad spectrum medicines used as a source of statin. A dose of 2.5mg/kg/day in rabbits was used to compare the efficiency of the test drug, EO, with the standard drug according to Ahmed *et al.*, (2015).

High Fat Diet: High fat diet (HFD) was made as described by Anjum *et al.*, (2014) to induce hypercholesterolemia by feeding a daily dose of 10 ml/kg/body weight orally as described by Ghuffar *et al.* (2014).

Study Plan: The experiment was performed for eight weeks. The animals were weighed, numbered and randomly separated into five groups with five animals per group as follows:

Group I served as a control group which received only Normal Saline. Group II was fed on normal diet and received a daily dose of 600mg/Kg of EO. Group III was fed on normal diet and a daily dose of HFD. Group IV received a daily dose of HFD and the test drug

(600mg/Kg of EO). Group V received a daily dose of HFD and Atorvastatin (2.5mg/kg/day).

Dose Administration: The powdered EO dose and Atorvastatin dose were measured and mixed with normal saline, while HFD was directly fed to the animals by an intra-gastric feeding tube as described by Dhuley (1999).

Serum Analysis: Feeding and maintenance of experimental animals was monitored very carefully until the end of the study. After eight weeks, 2ml blood was collected from marginal ear vein with the help of a capillary tube from each group to quantify the various parameters of lipid profile. Animals were fasted for at least 14 hours before the blood collection. The collected blood samples were centrifuged at 2500 x g at 20 °C for 10 minutes and serum thus obtained was stored at -20 °C as described by Anjum *et al.*, (2014) and Yaqub *et al.*, (2018). Serum was analyzed by calorimetric test for quantitative estimation of total cholesterol (TC), LDL, HDL, and Triglycerides (TG).

Estimation of Antioxidants: The rabbits were sacrificed and liver and kidneys were separated after dissection for the estimation of antioxidants. Tissues were homogenized in Phosphate-buffered saline (PBS) 7.4 pH and homogenate was sonicated and centrifuged at 20000 g for 15 min at 4 °C temperature. The supernatant was divided into two parts; one aliquot was stored at -20 °C while the other was used for estimation of antioxidants. Superoxide dismutase, Catalase and Glutathione peroxidase activity was measured as described by Sun *et al.*, (1988), Kawamura *et al.*, (1991), Paglia and Valentine (1967), respectively.

Statistical Analysis: Kolmogorov Smirnov's test was applied to evaluate the normal distribution of data. Data were presented as mean \pm S.D. The data were analyzed by one-way analysis of variance (ANOVA), followed by Dunnett's comparison tests using statistical package SPSS (Version 13.0 SPSS Inc., Chicago, IL, USA). Results with $P < 0.05$ were considered statistically significant.

Table 1. Comparative lipid profile of different experimental groups and effects of EO powder and atorvastatin on lipid profile after eight weeks of experimentation.

Animal Groups	Total Cholesterol (mg/dl)	LDL (mg/dl)	HDL (mg/dl)	Triglycerides (mg/dl)	Atherogenic Index
Group 1	105.79 \pm 3.37	63.32 \pm 2.34	34.19 \pm 2.45	72.54 \pm 1.98	1.80 \pm 0.12
Group 2	98.61 ^a \pm 1.29	44.28 ^a \pm 1.33	45.86 ^a \pm 1.40	61.28 ^a \pm 2.55	0.63 ^a \pm 0.07
Group 3	271.38 ^a \pm 5.88	287.30 ^a \pm 4.65	17.93 ^a \pm 1.84	227.65 ^a \pm 3.79	17.50 ^a \pm 1.20
Group 4	127.47 ^b \pm 3.24	67.72 ^b \pm 1.66	29.28 ^b \pm 1.39	88.91 ^b \pm 3.12	2.54 ^b \pm 0.03
Group 5	91.48 ^c \pm 3.57	39.59 ^c \pm 1.23	54.49 ^c \pm 1.33	59.76 ^c \pm 2.19	1.68 ^c \pm 0.34

^a: $P < 0.05$ ^a when compared the normal group to group II & III.

^b: $P < 0.05$ ^b when compared the normal group to hyperlipidemic and control group

^c: $P < 0.05$ ^c when compared the statin group to the normal group

RESULTS

A significant increase in all the lipid parameters, except for HDL was observed following the administration of HFD when compared to control (Table 1). Mean TC value amounted to 105.79 \pm 3.37 mg/dl in control rabbits, whereas in fat-fed rabbits this value amounted to 271.38 \pm 5.88mg/dl after eight weeks of the experimentation period. In the control group, mean values of LDL and TG were 63.32 \pm 2.34 mg/dl and 72.54 \pm 1.98 mg/dl, respectively; however, a significant increase was noted in both of these parameters in the HFD-fed animal group. Both LDL and TG levels decreased significantly ($P < 0.05$) due to administration of the test drug. The TG in the HFD-fed group was significantly higher ($P < 0.05$) as compared to the control group (72.54 \pm 1.98 mg/dl). Triglycerides level also decreased almost to the standard drug treatment level due to EO treatment. HDL decreased significantly (17.93 \pm 1.84mg/dl) with the increased HFD intake. HDL level was observed higher in rabbits when they were administered EO with normal feed. EO maintained the HDL at significant (29.280 \pm 1.39 mg/dl) high level even when administered with HFD as shown in Figure 1a. High fat diet led to significant decrease ($P < 0.05$) of SOD, CAT and GSH in liver and kidney of HFD group as shown in Table 2. *Emblica officinalis* powder significantly increased the content of antioxidant enzymes in liver and kidney tissue. Results also suggest that EO has better antioxidative potential than standard drug, statin; however, it was also noted that statin also relieved the oxidative stress of liver and kidney tissues in HFD group. The study revealed that concomitant administration of the EO powder at a dose of 600mg/kg body weight along with HFD in the study animals resulted insignificant decrease in the concentration of all antioxidant enzymes and lipid parameters except for HDL. The level of HDL increased ($P < 0.05$) in the EO powder fed rabbits. Moreover, the chances for atherosclerosis also increased with the HFD as shown in Figure 1b. *Emblica officinalis* powder and atorvastatin decrease the atherogenic index and atherosclerosis.

Table 2. Concentration of antioxidant enzymes in liver and kidneys of various experimental groups after eight weeks of experimentation

Animal Groups	SOD(mg ⁻¹)		CAT(mg ⁻¹)		GSH(m mol 100 g ⁻¹)	
	Liver	Kidney	Liver	Kidney	Liver	Kidney
Group 1	10.24±0.78	17.08±0.23	73.43±0.82	35.16±0.92	512±3.46	246±5.81
Group 2	10.89±0.12	18.93±0.31	75.21±0.34	37.29±0.29	529±7.21	252±1.19
Group 3	06.59 ^a ±0.67	10.03 ^a ±0.19	38.73 ^a ±0.95	20.45 ^a ±0.35	442 ^a ±8.36	211 ^a ±6.67
Group 4	10.40 ^b ±0.47	17.93 ^b ±0.27	74.18 ^b ±0.49	37.14 ^b ±0.28	522 ^b ±9.15	249 ^b ±2.32
Group 5	08.99 ^c ±0.76	16.28 ^c ±0.91	64.34 ^c ±0.03	33.46 ^c ±0.18	508 ^c ±0.22	241 ^c ±0.47

^a: P<0.05 ^a when compared the hyperlipidemic group to the normal group;

^b: P<0.05 ^b when compared the EO group to hyperlipidemic group;

^c: P<0.05 ^c when compared the statin group to the normal group.




Figure 1a: Mean serum lipid parameters in five groups of experimental rabbits receiving different treatments at the end of 8 week of experimentation.




Figure 1b: Mean atherogenic index in five groups of experimental rabbits receiving different treatments at the end of 8 weeks of experimentation.

DISCUSSION

Rossouw (2015) reported similar findings as observed in the present study, indicating that the intake of HFD is directly proportional to the cholesterol level in the circulating blood. Further, free radicals are produced by HFD leading to hyperlipidaemia and oxidative stress. Lipid peroxidation of high fats leads to high serum cholesterol levels (Anjum *et al.*, 2014). According to Shyamala *et al.* (2003) the abnormal lipid profile is created by the free radicals produced by the HFD. These free radicals create an oxidative stress leading to the abnormal lipid profile as observed in the current study. It is considered that high TG and TC levels were due to these free radicals. Oxidative stress leads to lipid peroxidation which is the one cause of many diseases, such as diabetes mellitus and coronary heart diseases. Higher LDL and TG levels are the core cause of atherosclerosis, insulin resistance and glucose resistance (Ghuffar *et al.*, 2014).

Administration of EO powder nearly maintained the lipid profile. The anti-hyperlipidemic effects of EO powder are quite comparable with the standard drug (atorvastatin). EO powder may have achieved the results because it is already known that it possesses anti-lipo-oxidative properties (Kumar *et al.*, 2012b). This powder contains various chemicals like, flavonoids, poly-phenolic compounds, and dietary fibres, etc., all of which have been reported as anti-hyperlipidemic compounds. The mechanism as to how the level of TG, TC and LDL decreased is not clear but several studies indicate that EO is rich in flavonoids and polyphenols which are natural antioxidants (Ghuffar *et al.*, 2014; Sairam *et al.*, 2002). These natural antioxidants perhaps stop the lipid peroxidation of fats resulting in the decreased levels of LDL, TG and total cholesterol in the blood (Scartezini *et al.*, 2006).

Previously, it has been proven that high fat intake significantly alters the antioxidative defence mechanism against lipid peroxidation in animals (Çelik *et al.*, 2017). It has been reported that hyper-lipidemia blocks the antioxidant defence system by reducing the activity of catalase and SOD which in turn leads to increased lipid peroxide content thus consequently toxic intermediates are produced. Furthermore, HDL results in the decreased concentrations of tissue glutathione level (Lennicke *et al.*, 2017). When experimental rabbits were treated with EO powder containing flavonoids, the glutathione levels were improved. Rutin (natural flavonoids) raised the level of glutathione activity in the tissues of mice when gastric lesions were induced in them (Ganeshpurkar and Saluja 2017). Studies reveal that flavonoids stimulate antioxidative enzymes to cope the oxidative stress (Semiz *et al.*, 2017).

Higher intake of fats creates an oxidative stress inside the body which triggers several

pathways/mechanisms for the lipid peroxidation. Cu²⁺-induced LDL oxidation, membrane lipid peroxidation and ADPFe³⁺/NADPH-induced lipid peroxidation are the most common pathways which are altered by the oxidative stress to create hyperlipidaemia. Antioxidant compounds like ascorbic acid, emblicanins, corilagin, ellagic acid and methyl gallate, etc., present in EO powder inhibit these pathways and directly regulate the abnormal lipid production (Asgary *et al.*, 2013). Ascorbic acid and other antioxidant compounds present in EO are also responsible for downregulation of 3 hydroxy 3 methyl-glutaryl coenzyme A reductase activity resulting in the increased biliary cholesterol excretion thus inhibiting the cholesterol absorption through intestine (Hirose *et al.*, 1991). It is also considered that EO powder reduces the high lipids in the body by producing some antioxidant enzymes like glutathione peroxidase, superoxide dismutase and catalases etc., as these have been reported to be produced by other herbs (*Cinnamomum verum*, *Syzygium aromaticum* and *Amoma subulatum*) when administered to HFD rats (Dhuley, 1999; Shyamala *et al.*, 2003). Since the antioxidant ingredients from plants are helpful against tissue damage, heart disease and cancer, and have now attracted the attention of scientific community and food manufacturers, therefore, the future belongs to functional foods that have beneficial health impacts (Coman *et al.*, 2017).

REFERENCES

- Abbasi, A. M., M. Khan, M. Ahmad, M. Zafar, S. Jahan and S. Sultana (2010). Ethnopharmacological application of medicinal plants to cure skin diseases and in folk cosmetics among the tribal communities of North-West Frontier Province, Pakistan. *J. Ethnopharmacol.* 128(2): 322-335.
- Ahmad, I., Z. Mehmood and F. Mohammad (1998). Screening of some Indian medicinal plants for their antimicrobial properties. *J. Ethnopharmacol.* 62(2): 183-193.
- Ahmed, S., S. Gul, M. Zia-Ul-Haq and M. Riaz (2015). Hypolipidemic effects of nimesulide and celecoxib in experimentally-induced hypercholesterolemia in rabbits. *Turk. J. Med. Sci.* 45(2):277-83.
- Anjum, K.M, U. Sayyed, A. Ullah, M. S. Mughal, A. Yaqub, M. A. Rashid and M. Z. Yousaf (2014). Anti-hypercholesterolemic and anti-atherogenic activity of *Terminalia chebula* fruit in normal and cholesterol fed rabbits. *The J. Anim. Plant Sci.* 24(6): 1618-1622.
- Asgary, S., M. Rafieian-Kopaei, S. Najafi, E. Heidarian and A. Sahebkar (2013). Antihyperlipidemic effects of *Sesamum indicum* L. in rabbits fed a high-fat diet. *Sci. World J.* 10(2): 1-5.

- Bazmandegan, G., M. T. Boroushaki, A. Shamsizadeh, F. Ayoobi, E. Hakimizadeh and M. Allahtavakoli (2017). Brown propolis attenuates cerebral ischemia-induced oxidative damage via affecting antioxidant enzyme system in mice. *Biomed. Pharmacother.* 85: 503-510.
- Binti Anzian, A., R. Sukor, N. Saari, C. W. N. S. binti Che and W. Sapawi (2017). Chemical composition and antioxidant activity of Torch Ginger (*Etlingera elatior*) flower extract. *Food Appl. Biosci. J.* 5(1): 32-48.
- Cappelli-Bigazzi, M., S. Rubattu, C. Battaglia, R. Russo, I. Enea, G. Ambrosio, M. Chiariello and M. Volpe (1997). Effects of high-cholesterol and atherogenic diets on vascular relaxation in spontaneously hypertensive rats. *Am. J. Physiol. Heart Circ. Physiol.* 273(2): H647-H654.
- Çelik, N., A. Vurmaz and A. Kahraman (2017). Protective effect of quercetin on homocysteine-induced oxidative stress. *Nut.* 33: 291-296.
- Coman, M., M. Verdenelli, S. Silvi, C. Cecchini, R. Gabbianelli, E. Amadio, C. Orpianesi and A. Cresci (2017). Knowledge and acceptance of functional foods: a preliminary study on influence of a synbiotic fermented milk on athlete health. *Intl. J. Probio. Prebio.* 12(1): 27-35.
- Dhuley, J. (1999). Anti-oxidant effects of cinnamon (*Cinnamomum verum*) bark and greater cardamom (*Amomum subulatum*) seeds in rats fed high fat diet. *Indian J. Exp. Biol.* 37: 238-242.
- Ganeshpurkar, A. and A. K. Saluja (2017). The pharmacological potential of rutin. *Saudi Pharm. J.* 25(2): 149-164.
- Ghuffar, A., T. Ahmad and M. N. Mushtaq (2014). Antihyperlipidemic effect of *Berberis orthobotrys* in hyperlipidemic animal models. *Bangladesh J. Pharmacol.* 9(3): 377-382.
- Goldstein, J. L. and M. S. Brown (2015). A Century of Cholesterol and Coronaries: From Plaques to Genes to Statins. *Cell* 161(1): 161-172.
- Herron, K. L., I. E. Lofgren, M. Sharman, J. S. Volek and M. L. Fernandez (2004). High intake of cholesterol results in less atherogenic low-density lipoprotein particles in men and women independent of response classification. *Metabol.* 53(6): 823-830.
- Hirose, N., T. Inoue, K. Nishihara, M. Sugano, K. Akimoto, S. Shimizu and H. Yamada (1991). Inhibition of cholesterol absorption and synthesis in rats by sesamin. *J. Lipid Res.* 32(4): 629-638.
- Illingworth, D. R. (1987). Lipid-lowering drugs. *Drugs* 33(3): 259-279.
- Jain, P., H. Sharma, F. Basri, B. Baraik, S. Kumari and C. Pathak (2014). Pharmacological Profiles of Ethno-Medicinal Plant: *Plumbago Zeylanica* L.-A Review. *Int. J. Pharm. Sci. Rev. Res.* 24(1): 157-163.
- Jain, R., R. Pandey, R. Mahant and D. Rathore (2015). A review on medicinal importance of *Emblca officinalis*. *Int. J. Pharm. Sci. Res.* 6(1): 72-76.
- Kawamura, T., T. Yoshioka, T. Bills, A. Fogo and I. Ichikawa (1991). Glucocorticoid activates glomerular antioxidant enzymes and protects glomeruli from oxidant injuries. *Kidney Int.* 40(2): 291-301.
- Kawasumi, K., N. Kashiwado, Y. Okada, M. Sawamura, Y. Sasaki, E. Iwazaki, N. Mori, I. Yamamoto and T. Arai (2014). Age effects on plasma cholesterol and triglyceride profiles and metabolite concentrations in dogs. *Vet. Res.* 10(1): 57.
- Khan, K. (2009). Roles of *Emblca officinalis* in medicine-a review. *Bot. Res. Int* 2(4): 218-228.
- Kumar, A., A. Singh and J. Dora (2012a). Essentials perspectives for *Emblca officinalis*. *Int. J. Pharm. Chem. Sci.* 1(1): 11-18.
- Kumar, K. S., D. Bhowmik, A. Dutta, A. P. Yadav, S. Paswan, S. Srivastava and L. Deb (2012b). Recent trends in potential traditional Indian herbs *Emblca officinalis* and its medicinal importance. *J. Pharm. Phytochem.* 1(1): 18-28.
- Kureishi, Y., Z. Luo, I. Shiojima, A. Bialik, D. Fulton, D. J. Lefer, W. C. Sessa and K. Walsh (2000). The HMG-CoA reductase inhibitor simvastatin activates the protein kinase Akt and promotes angiogenesis in normocholesterolemic animals. *Nature Med.* 6(9): 1004-1010.
- Lenicke, C., J. Rahn, A. P. Kipp, B. P. Dojčinović, A. S. Müller, L. A. Wessjohann, R. Lichtenfels and B. Seliger (2017). Individual effects of different selenocompounds on the hepatic proteome and energy metabolism of mice. *Biochim. Biophys. Acta.* 1861(1): 3323-3334.
- Paglia, D. E. and W. N. Valentine (1967). Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *Trans. Res.* 70(1): 158-169.
- Pahan, K. (2006). Lipid-lowering drugs. *Cell. Mol. Life Sci.* 63(10): 1165-1178.
- Qasim, M., Z. Abideen, M. Adnan, S. Gulzar, B. Gul, M. Rasheed and M. Khan (2017). Antioxidant properties, phenolic composition, bioactive compounds and nutritive value of medicinal halophytes commonly used as herbal teas. *South African J. Bot.* 110: 240-250.
- Rossouw, J. (2015). Serum cholesterol as a risk factor for coronary heart disease revisited. *South African J. Clin. Nut.* 28(1): 34-37.

- Sairam, K., C. V. Rao, M. D. Babu, K. V. Kumar, V. Agrawal and R. K. Goel (2002). Antiulcerogenic effect of methanolic extract of *Emblica officinalis*: an experimental study. *J. Ethnopharmacol.* 82(1): 1-9.
- Scartezzini, P., F. Antognoni, M. Raggi, F. Poli and C. Sabbioni (2006). Vitamin C content and antioxidant activity of the fruit and of the Ayurvedic preparation of *Emblica officinalis* Gaertn. *J. Ethnopharmacol.* 104(1): 113-118.
- Semiz, A., G. C. Turgut, S. Karakurt, H. Akca, S. Arslan, O. Adali and A. Sen (2017). In vivo examination of the effects of hydroxycinnamic acid on xenobiotic metabolizing and antioxidant enzymes. *Arch. Biol. Sci.* 69(1): 103-110.
- Shyamala, M., M. Venukumar and M. Latha (2003). Antioxidant potential of the *Syzygium aromaticum* (Gaertn.) Linn.(Cloves) in rats fed with high fat diet. *Indian J. Pharmacol.* 35(2): 99-103.
- Sun, Y., L. W. Oberley and Y. Li (1988). A simple method for clinical assay of superoxide dismutase. *Clin. Chem.* 34(3): 497-500.
- Varbo, A., M. Benn and B. G. Nordestgaard (2014). Remnant cholesterol as a cause of ischemic heart disease: evidence, definition, measurement, atherogenicity, high risk patients, and present and future treatment. *Pharmacol. Ther.* 141(3): 358-367.
- Yaqub, A., K. M. Anjum, A. Munir and H. M. W. A. Khan (2018). Evaluation of acute toxicity and effects of sub-acute concentrations of copper oxide nanoparticles (CuO-NPs) on hematology, selected enzymes and histopathology of liver and kidney in *Mus musculus*. *Indian J. Anim. Res.* 52(1): 92-98. DOI: 10.18805/ijar.v0iOF.8489.