

**PREDICTING THE POTENTIAL HABITAT AND DISTRIBUTION OF
WESTERN TRAGOPAN (*TRAGOPAN MELANOCEPHALUS*) IN SELECTED AREAS OF
AJ&K, PAKISTAN: A MAXENT MODELING APPROACH**

H. Ali^{1,2*}, U. Akram¹, S. Abbas¹, M. S. Ahmed², F. M. Qamer³, B. Khan¹, M. N. Awan⁴, Z. Ali², A. A. Chaudhry¹,
A. Saghir² and M. Nagai⁵

¹World Wide Fund for Nature (WWF) – Pakistan

²Zoology Department, University of the Punjab, Lahore, Pakistan

³International Centre for Integrated Mountain Development (ICIMOD), Nepal

⁴Himalayan Nature Conservation Foundation, AJ&K, Pakistan

⁵Geographic Information Centre, AIT, Thailand

*Corresponding author Email: sukheraji@gmail.com

ABSTRACT

Western Tragopan (*Tragopan melanocephalus*) is a beautiful pheasant species and is endemic to western Himalayas. It has large distribution range but different populations are fragmented within the western Himalayan moist temperate forests. Its distribution in Pakistan is restricted to some parts of Khyber Pakhtunkhwa and Azad Jammu & Kashmir (AJ&K). This species is globally 'vulnerable' since its fragmented small populations are declining due to continuing forest loss and habitat destruction throughout its range. This study was carried out to model the potential distribution and habitat of Western Tragopan in Neelum and Muzaffarabad districts of AJ&K by incorporating the physical, biological and climatic variables into MaxEnt model. L and cover data, Digital Elevation Model (DEM), slope, aspect, Normalized Difference Vegetation Index (NDVI), temperature, precipitation, topographic and sighting datasets were processed in Geographic Information System (GIS) to produce meaningful predictor variables, subsequently used in the MaxEnt software. The results showed that 11,112 ha area (2% of the total study area) is highly suitable for Western Tragopan, 30,248 ha area (5.3%) moderately suitable while 525,445 ha (92.7%) area is unsuitable. Jackknife test evaluated the importance of predictor variables and DEM, precipitation and land cover were found to be the most important variables for this study. The study evaluated the species-habitat relationship within the study area and can be helpful in the management and *in-situ* conservation planning of the species in its distribution range.

Key words: Western Tragopan, Potential distribution, MaxEnt modeling, *in-situ* conservation

INTRODUCTION

Western Tragopan (*Tragopan melanocephalus*) is a beautiful pheasant species and is endemic to western Himalayas (Bird Life International, 2015). It is a shy bird, usually occurs singly, roost on trees but feeds on the ground in early morning and late afternoon (Roberts, 1991). Western Tragopan is a low density and habitat specific species, prefers extremely steep terrain and is closely associated with open moist deciduous (e.g. maple, oak, birch and walnut) and/or coniferous temperate vegetation (especially silver fir, blue pine and spruce) with dense shrubby undergrowth. It can easily survive in extreme cold at high altitudes. Although, it forages close to the upper margins of the forest (i.e. 3,350m) but in winter it descends as low as 2,150m or up to snowline (Roberts, 1991; Whale, 1996; Anwar *et al.*, 2006; Bird Life International, 2015).

The Western Tragopan is the most westerly distributed among the five Tragopan species occurring in India and Pakistan in the Himalayas. In Pakistan its distribution is patchy and small population survives in

Duber and Palas valleys in Indus Kohistan, Kaghan valley, Pir Hasimar, Machiara, Pir Chinasi and Salkhala in Azad Kashmir. (Roberts, 1991; Whale, 1996; Awan, 2010). This species is globally 'vulnerable' since its fragmented small populations are declining throughout its range. (Roberts, 1991; BirdLife International 2015). The reasons for the population decline are attributed to the habitat loss due to forest degradation, hunting and trapping (Whale, 1996; Ashraf *et al.*, 2004).

There have been very few studies to evaluate the distribution and status of Western Tragopan in Pakistan (Grimmett and Craig, 1984; Roberts, 1991; Bean *et al.* 1994; Whale 1996; Ghafoor & Nawaz, 1999; Awan, 2010). All these studies were primarily conducted through ground-based field surveys. Ashraf *et al.* (2004) evaluated the habitat and distribution of Western Tragopan in Palas Valley using a combination of Landsat based land cover, terrain data and field survey data. Although traditional survey techniques are important in assessing the distribution and habitat preference of the species but the development of geospatial technologies has provided a complimentary system to ground-based

wildlife studies (Ali, *et al*, 2010; Qamer *et al*, 2011; Matin *et al*, 2012).

With the development of new statistical and geospatial techniques, the development of predictive species distribution models (SDMs) has increased in ecology, biogeography and conservation biology (Guisan and Zimmerman, 2000; Guisan and Thuiller, 2005). Currently a wide range of species distribution models are being used worldwide (Stockwell and Peters, 1999; Friedman *et al*, 2000; Breiman, 2001; Wikle, 2003; Philips *et al*, 2006). These SDMs mainly use species occurrence data with ecological/environmental variables (vegetation, temperature, precipitation etc.) and topographic variables (elevation model, slope, aspect etc.) to create a distribution and habitat model of a species (Guisan and Thuiller, 2005; Elith *et al*, 2006; Zimmermann *et al*, 2010)

In this study we used MaxEnt model to study the potential distribution and habitat of Western Tragopan in selected areas of AJ&K Pakistan, based on the geographic distribution data and remote sensing/GIS derived predictor variables. MaxEnt is a predictive species distribution model which is based on the principle of maximum entropy and recognized as the most useful

model compared to the other SDMs (Elith *et al*, 2006; Phillips *et al*, 2006). Key objectives of the study were 1) to predict the potential geographic range of the species within the study area and 2) to identify the environmental/geographical variables closely associated with the species influencing its geographical distribution.

MATERIALS AND METHODS

Study Area: The study area includes Muzaffarabad and Neelum Districts of Azad Jammu & Kashmir located in the north eastern part of Pakistan (Fig. 1). The study area falls in the moist temperate zone of western Himalayas with extreme winters and snowfall. The mean annual rainfall is 1526.7 mm, with 84.5 rainy days per year. The rainiest month is July with a mean rainfall of 327.6 mm, while the driest month is November receiving a mean rainfall of only 35.4 mm (Anonymous, 2005). The study area is part of the Western Himalayan Ecoregion that is included in 200 important global ecoregions ranked by WWF based on their conservation status (Wikramanayake *et al*, 2002).




Figure 1: Study area map

Land cover mapping: For the development of land cover of the study area, ALOS multispectral satellite images with 10m spectral resolution were used. Field surveys were carried out to collect ground truth data using Garmin Map 76 CSX GPS receiver with ± 5 m accuracy. A total of 300 GPS-linked data descriptive

points were collected during the surveys. Object Based Image Analysis (OBIA) was performed using Definien Developer 7.0 software based on the field survey data and a land cover map of the study area was developed (Fig. 2).




Figure 2: Map showing species sighting data and land cover of the study area

Species sighting data and Ecological/Environmental Variables: Thirty two species sighting points were collected from the unpublished survey reports of AJ&K Wildlife Department (Fig. 2). Ten environmental variables were used for the study viz., land cover derived from ALOS satellite data, 30m resolution digital elevation model (DEM) downloaded from ASTER GDEM (<http://gdem.ersdac.jspacesystems.or.jp/>), aspect and slope (derived from DEM), temperature & precipitation downloaded from WorldClim datasets (<http://www.worldclim.org/>), normalized difference vegetation index (NDVI) from Landsat and raster layers of distance to drainage, distance to ridges, distance to roads, derived from topographic data.

Spatial modeling: Maximum Entropy (MaxEnt) modelling software (version 3.3.3e) was used to predict the potential distribution and habitat of the Western Tragopan in the study area. Maxent modeling is a general-purpose method which uses presence-only data and makes predictions of species' potential distribution/habitat from incomplete information and its origin lie in statistical process of maximum entropy (Jaynes, 1963; Phillips *et al.*, 2006). MaxEnt uses the species sighting data with other ecological/environmental/geographical variables and calculates the likelihood of presence of a given species or

number of species within the study area (Phillips *et al.*, 2006; Phillips and Dudik, 2008).

ArcGIS 9.3 and ENVI 4.5 software were used to process and develop the standardized spatial data layers of DEM, slope, aspect, land cover, distance to drainage, distance to ridges, and distance to roads, precipitation and temperature to be used in MaxEnt modeling software. 70% of species sighting data was used as training data and 30% was used as test data. Final potential distribution and habitat map of Western Tragopan was produced with a range of values from 0 to 1 which were regrouped in to three different classes viz., unsuitable, moderately suitable and highly suitable.

RESULTS AND DISCUSSION

MaxEnt model produced a continuous prediction with values ranging from 0 to 1. The values are illustrated here using blue color (0-0.5) representing unsuitable habitat for the species, green color (0.51-0.7) representing moderately suitable habitat while red color (0.71-1) which represents highly suitable habitat for Western Tragopan within the study area (Fig. 3). The results of the study revealed that 11,112 ha area was highly suitable which is 2% of the total study area, 30,248 ha (5.3%) area

was moderately suitable while 525,445 ha (92.7%) area is unsuitable.




Figure 3: Predicted potential habitat/distribution of Western Tragopan in the study area

The jackknife regularized training gain results showed DEM, precipitation and land cover, as main variables influencing the Western Tragopan distribution (Fig. 4). Other important variables include NDVI, distance to roads and aspect. This test actually determines

the contribution of each variable by testing the impact of the variable removing it from the model and also testing the impact of each variable by running the model only with that variable (Philips *et al*, 2006)




Figure 4. The Jackknife test output showing the relative importance of predictor variables used in the study

The MaxEnt model output provided adequate results with the given set of training/test data and the model fitness as measured by area under receiver operating characteristics (ROC) curve (AUC) was with a high value of 0.952 for the training data and 0.871 for the

test data (Fig. 5). This showed better ability of the MaxEnt model to discriminate between suitable and unsuitable areas for the Western Tragopan occurrence. The ROC/AUC analysis has been extensively used in species habitat and distribution modeling and is

recognized as the best measure of model performance (Philips *et al.*, 2006; Elith *et al.*, 2011).




Figure 5: The ROC curve for the training and test data

Current study predicted the habitat and suitable distribution areas of Western Tragopan in selected areas of its distribution range using SDM tool and for the first time quantified the habitat of the study species in the study area. Earlier, Ashraf *et al.*, (2004) conducted a GIS based overlay analysis to map the habitat and distribution of Western Tragopan in Palas Valley, Indus Kohistan, and Pakistan. The areas which are predicted suitable for Western Tragopan but not occupied currently can be considered for introduction/reintroduction of this species and future field efforts should be focused more on the identified potential habitats to assess the conservation status of this species.

This study is a pioneer effort in habitat suitability mapping for the Western Tragopan in Pakistan and can be of great help in developing effective management strategies for the *in-situ* conservation of this regionally endemic bird within the study area. Further, current predicted species distribution and habitat model developed with MaxEnt can be used with the global climate change scenarios to forecast the potential changes in distribution range.

Acknowledgements: The study was funded by Japan Aerospace Exploration Agency (JAXA) with technical assistance from Geo-Informatics Center, Asian Institute of Technology (GIC-AIT), Bangkok Thailand. We are also thankful to Department of Wildlife and Fisheries, Govt. of Azad Jammu and Kashmir for providing assistance in data collection during the field surveys.

REFERENCES

- Ali, H., R. Saleem, F.M. Qamer, W. A. Khan, S. Abbas, K. Gunasekara, M. Hazarika, M.S. Ahmed, and M. Akhtar (2010). Habitat Evaluation of Smooth-coated Otter (*Lutrogale perspicillata*) in Indus Plains of Pakistan using Remote Sensing and GIS. *Int Arch Photogramm, Remote Sensing Spatial Inform Sci.*, 38(8): 127 - 132.
- Anonymous. (2005). Revised Management Plan of Machiara National Park. Department of Wildlife and Fisheries. Government of Azad Jammu and Kashmir. Muzaffarabad.
- Anwar, M., Q. Zaman, and R. A. Minhas (2006). Study Report on Population of Selected Wildlife Species in Machiara National Park, AJK. Protected Areas Management Project, Muzaffarabad.
- Ashraf, S, R. Nawaz, A. Daud and F. M. Qamer (2004). Mapping Western Tragopan *Tragopan melanocephalus* habitat and distribution pattern in the Palas Valley, Pakistan using landcover, terrain and field survey data. *Proc. of Int. Sym on Galliformes*, Deradun, India.
- Awan, M. N. (2010). Status and conservation of Western Tragopan Pheasant in and around Salkhala Game Reserve, District Neelum, Azad Kashmir, Pakistan. Final Progress Report Submitted to Oriental Bird Club. UK.
- Bean, N. J., P.J. Beanstead, D. Showler and P.A. Whittington (1994). Survey of Western Tragopan (*Tragopan melanocephalus*) in the Palas Valley, NWFP – Spring 1994. Final report submitted to Himalayan Jungle Project,

- Islamabad, Pakistan and BirdLife International, Cambridge, UK.
- Bird Life International (2015), Species factsheet: *Tragopan melanocephalus*. Downloaded from <http://www.birdlife.org> on 08/03/2015
- Breiman, L. (2001). Random forests. *Machine Learning*, 45: 5–32.
- Elith, J., C. H. Graham, R. P. Anderson, M. Dudik, S. Ferrier, A. Guisan, R. J. Hijmans, F. Huettmann, J. R. Leathwick, A. Lehmann, J. Li, L.G. Lohmann, B.A. Loiselle, G. Manion, C. Moritz, M. Nakamura, Y. Nakazawa, J. M. Overton, A. T. Peterson, S.J. Phillips, K. Richardson, R. Scachetti Pereira, R. E. Schapire, J. Soberon, S. Williams, M. S. Wisz, and N. E. Zimmermann (2006). Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, 29: 129–151.
- Elith, J., S. J. Phillips, T. Hastie, M. Dudik, Y. E. Chee, C. J. Yates (2011). A statistical explanation of MaxEnt for ecologists. *Divers. Distrib.* 17: 43–57.
- Friedman, J. H., T. Hastie, and R. Tibshirani (2000). Additive logistic regression: a statistical view of boosting. *Ann. Stat.*, 28: 337–407.
- Ghafoor, A. and R. Nawaz (1999). Survey of western tragopan and other pheasant species in Bar Palas, Kohistan. Unpublished report of WWF-Pakistan/Himalayan Jungle Project.
- Grimmett, R. and R. Craig (1984). Report on survey of the Western Tragopan in Pakistan (unpublished), submitted to National Council for Conservation of Wildlife, Islamabad.
- Guisan, A., and N. E. Zimmermann (2000). Predictive habitat distribution models in ecology. *Ecological Modeling*, 135: 147–186.
- Guisan, A., and W. Thuiller (2005). Predicting species distribution: offering more than simple habitat models. *Ecology Letters*, 8:993–1009.
- Jaynes, E. T. (1963). Information Theory and Statistical Mechanics. In Ford, K. (ed.). *Statistical Physics*. New York: Benjamin. 181.
- Matin, S., V. S. Chitale, M. D. Behera, B. Mishra, P. S. Roy (2012), Fauna data integration and species distribution modeling as two major advantages of geoinformatics-based phytobiodiversity study in today's fast changing climate. *Biodivers. Conserv.* 21: 1229–1250.
- Phillips, S. J., and M. Dudik (2008). Modeling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography*, 31: 161–75.
- Phillips, S. J., R. P. Anderson and R. E. Schapire (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modeling*, 190: 231–259.
- Qamer, F. M., H. Ali, S. Ashraf, A. Daud, H. Gillani, H. Mirza, H.U. Rehman (2011). Distribution and Habitat Mapping of Key Fauna Species in Selected Areas of Western Himalaya, Pakistan. *J. of Anim and Plant Sci.*, 21(2 Suppl.): 396 – 399
- Roberts, T. J. (1991). *The Birds of Pakistan, Volume 1. Non-passeriformes*. Oxford University Press, Karachi.
- Stockwell, D. and D. Peters (1999). The GARP modeling system: problems and solutions to automated spatial prediction. *Int. J. Geogr. Inform. Sci.*, 13: 143–158.
- Whale, R. (1996). Survey Report of Western Tragopan in the Palas Valley (unpublished), Himalayan Jungle Project, Abbottabad, Pakistan.
- Wikle, C. K. (2003). Hierarchical Bayesian models for predicting the spread of ecological processes. *Ecology*, 84: 1382–1394.
- Wikramanayake, E., E. Dinerstein, C. Loucks, D. Olson, J. Morrison, J. Lamoreux, M. McKnight, P. Hedao (2002). *Terrestrial Ecoregions of the Indo-Pacific: A Conservation Assessment*. Island Press, Washington, DC.
- Zimmermann N. E., T. C. Edwards, C. H. Graham, P. B. Pearman and J. C. Svenning (2010). New trends in species distribution modeling. *Ecography*, 33:985–989.