

## **BIOLOGICAL EVALUATION, CHARACTERIZATION AND DISTRIBUTION OF SOME LICHENS OF HIMALAYAN REGION, PAKISTAN**

A. Ullah<sup>1</sup>, K. William<sup>3\*</sup>, A. Rehman<sup>2</sup>, S. W. Khan<sup>4</sup> and Z. Iqbal<sup>1</sup>

<sup>1</sup>Hazara University, Mansehra, Pakistan<sup>1</sup>,

<sup>2</sup>Institute of Natural and Management Sciences (INAM), Rawalpindi, Pakistan,

<sup>3</sup>Bioresource Research Centre (BRC), Islamabad, Pakistan

<sup>4</sup>Department of Biological Sciences, Karakoram International University, Gilgit, Pakistan.

\*Corresponding Author's Email: [kainaawatwill@hotmail.com](mailto:kainaawatwill@hotmail.com)

### **ABSTRACT**

Present study investigates the distribution, antibacterial, antioxidant, antitumor and cytotoxicity of the acetone and methanol extracts of ten lichen genera of *Dermetacarpon*, *Peltigera*, *Flavopunctelia*, *Flavoparmelia*, *Lecanora*, *Cladonia*, *Caloplaca*, *Collema*, *Rhizocarpon*, *Usnea* of Himalayan region (Mansehra) Pakistan. The bacterial strain used in this study was *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus bacillus*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Enterococcus faecalis*. The antibacterial activity of the lichen extract in both solvent was studied by Disc diffusion assay. The zone of inhibition of these lichen extracts were in range 7 to 12.7 mm in diameter. This study showed that *S. bacillus* is the most resistance and *S. epidermidis* is the most sensitive bacteria to the extract of all lichen genera. The antioxidant activity was investigated by 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging. The acetone extract in concentration of 1000ug/ml of *Collema* showed strongest DPPH radical scavenging activity while other acetone extracts also showed positive results. However, *Rhizocarpon* showed the least DPPH radical scavenging activity. Cytotoxic activity of different concentration of lichen extract was tested against Brine shrimp, *Artemia nauplii*. Result of cytotoxicity indicated that both methanol and acetone extract of *Flavopunctelia* were potent against brine shrimp. In potato disc assay, most of the acetone and methanol extracts of lichens showed significant (inhibition $\geq$ 20%) antitumor activity against *Agrobacterium tumefaciens*. Similarly, both acetone and methanol extracts of *Dermetocarpon* showed maximum (40%) tumor inhibition while *Collema* showed minimum tumor inhibition (16.66%).

**Key words:** Antibacterial activity, Antioxidant activity, Lichens, Himalayan region, Pakistan.

### **INTRODUCTION**

Lichen is a symbiotic association between algae and fungus. On the globe, they are represented by around 20,000 species (Shyam *et al.*, 2010), growing in terrestrial habitat on a variety of substrates i.e. bark, wood, leaves, rocks, soil and other fixture (Shah, 2011). They are important source of food for animals including man and are also used in alcohols, paints, perfume and pharmaceutical industries. Many natives of American, Indian and European tribes use it as their traditional medicine for treatments of many human and animal diseases (Karthikaidevi *et al.*, 2009). Lichens also produce various types of secondary metabolites i.e. phenolic compound, usnic acid, quinins, depsidones, deposesones and pulvinic acid derivatives etc. which are unique with respect to higher plants (Veranja *et al.*, 2005) having remarkable antibacterial, antiviral, antifungal, antioxidant and antitumor activities (Tatjana *et al.*, 2011). Concentrations usually 0.1-1% of dry thallus weight; sometimes up to 30% of these metabolites vary with the species and area in varying amounts. A number of studies are available on biochemical assays of these secondary metabolites of lichen (Santiago *et al.*, 2010; Marijana and

Rankovic, 2010; Krystle *et al.*, 2010) but lichens of Manshera or Pakistan are not studied yet.

### **MATERIALS AND METHODS**

**Collection of Lichen specimens:** Mansehra (Hazara Division) is North Eastern District (34°14' - 35° 11' NL; 72° 49' - 74° 08' E) of Khyber Pakhtoonkhawa, Pakistan covering an area of 4,579 km<sup>2</sup> (Ali, 2005). Lichen specimens were collected from Dader and Manroor valley, Lolosar and Behisal area of District Mansehra during 2013. The samples collected were *Dermetacarpon*, *Peltigera*, *Flavopunctelia*, *Flavoparmelia*, *Lecanora*, *Cladonia*, *Caloplaca*, *Collema*, *Rhizocarpon* and *Usnea*. All collected samples were sun/air dried by changing the papers at suitable intervals, mounted on a cardboard paper and covered with thick brown packet, numbered and labeled. Each specimen was then identified based upon morphology of the thallus and micro-chemical color tests (Smith, 1918: Chopta, 1934: Culberson and Kristinsson, 1970: Culberson, 1972: Shyam *et al.*, 2010: Aptroot and Iqbal, 2012).

**Distribution:** Samples of different lichen genera were collected from different localities of Manshera, Pakistan. *Dermetacarpon*, *Flavopuntelia*, *Collema* were collected from Dader, *Flavoparmelia*, *Cladonia* and *Peltigera* from Manoor Valley, *Lecanora* from Lalosar and *Caloplaca* from Behisal. The altitude of each location are given in Table 1.

**Preparation of Lichen extract:** The plant extracts of 10 lichen genera samples i.e. *Dermetacarpon*, *Peltigera*, *Flavopuntelia*, *Flavoparmelia*, *Lecanora*, *Cladonia*, *Caloplaca*, *Collema*, *Rhizocarpon* and *Usnea* were prepared using acetone and methanol. The weighed quantity one gram of each lichen was homogenized with 50 ml of respective solvents. Crude preparations were left overnight in orbital shaker at room temperature. It was then filtered and evaporated at 60°C. The extract was then weighed and then dissolved in distilled water to obtain final concentrations for each sample (Claudia *et al.*, 2014).

**Bacterial Strains:** Six bacterial strains (ATCC bacterial strains) (Gram +ve and -ve) were used in antibacterial assay i.e. *E.coli*, *Enterococcus faecalis*, *Staphylococcus bacillus*, *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Staphylococcus epidermidis*. These bacterial strains were cultured at 37 °C in 40ml of Liquid broth (LB) media and were maintained at 4 °C.

**Antibacterial Assay:** An antibacterial activity of each lichen extract was tested by measuring inhibition zone by disc diffusion assay (Sharma *et al.*, 2011). For this purpose Muller Hinton agar (bacteria) were seeded with appropriate inoculums. Paper discs (6 mm diameter) were impregnated with different concentrations of lichen extract and placed on organism seeded plates. Blank disc was impregnated with dimethyl sulphoxide (DMSO) and used as negative control while roxithromycin was used as positive control. The plates were incubated and the zones of inhibition around each disc were measured (Sati and Joshi, 2011)

**Cytotoxicity Assay:** Cytotoxicity of Lichen extracts was tested by Brine shrimp cytotoxicity assay, most widely used tool for detection of bioactive compound. A container filled with Brine solution was used for hatching of Brine shrimps eggs. Brine solution was prepared by dissolving 32 g of sea salt and 1.5% agar in distilled water. Ten shrimps were transferred to each tube containing brine solution 4.5ml/tube and different concentration of Lichen extract Acetone and Methanol (0.05 mg/ml, 0.01mg/ml and 0.001mg/ml) were added in each dilution for each sample to check its cytotoxicity. After 24 hours of incubation the number of surviving shrimps was counted by magnifying glass (Olowa and Muneza, 2013)

**Antitumor activity:** Antitumor activity was evaluated through Potato disc bioassay (Coker *et al.*, 2003). Potatoes taken from local market were surface sterilized with 20% bleach solution. The experiments were performed in two groups, extract of four lichen species and positive control. Stock solution of 0.05 mg/mL in DMSO was further diluted with sterilized water to achieve 0.01mg/mL and 0.001mg/mL concentration. Same concentration of positive control (Vincristine) was also prepared. Potato disc were made by cork borer and placed on 2% agar plates (6 disc/ plates). *Agrobacterium tumefaciens* culture of 48 hours was transferred on the surface of each plate (50µl). Same concentration (0.05mg/mL, 0.01mg/mL and 0.001mg/mL) of the lichen extract and positive control were applied to each plate. The disc were then stain with lugol's solutions (10% KI, 50% I) after 21 days of incubation at 28 °C. Number of tumor were counted under dissecting microscope and magnifying glass. Percentage inhibition was counted by following formula:

**Percentage inhibition:**  $100 - \frac{\text{No of tumor per sample}}{\text{Number tumor per control}} \times 100$ .

**Antioxidant activity:** The antioxidant activity of investigated lichen genera were reported through DPPH (1, 1- diphenyl-2 picrylhydrazyl) assay. Lichen extract of concentration 1000ug/ml, 500ug/ml, 250ug/ml and 125ug/ml prepared in methanol and acetone were used in this assay. Ascorbic acid was used as standard. (Aliyu *et al.*, 2012, Sharma and Kalikotay, 2012)

## RESULTS

The area of Manshera was covered in the current study and sample of *Dermetacarpon*, *Flavopuntelia*, *Collema* were collected from Dader, *Flavoparmelia*, *Cladonia* and *Peltigera* from Manoor Valley, *Lecanora* from Lalosar and *Caloplaca* from Behisal. The altitude of each location is given in Table 1. The antibacterial activities of the lichen extract were measured by Disc diffusion assay. The extract of these ten lichen genera showed variable range of zone of inhibitions against tested bacteria (Table 2). The maximum rate of antibacterial activity was observed in acetone extract of *Flavoparmelia* against *E.coli* (12.66±0.577) while methanol and acetone extract of *Peltigera* have no activity against *E.coli*, *S.bacillus* and *S.aureus*. Similar result was also recorded for *Flavopuntelia* and *Collema* against *S.bacillus*. The result of antioxidant activity of methanol and acetone extract and control is shown in Table 3 and 4. For methanol extract of concentration 1000ug/ml, the strongest (66%) DPPH radical scavenging activity was shown by the lichen genera *Cladonia*, followed by *Rhizocarpon* (63%) while in acetone maximum activity was *Rhizocarpon* (57%) and *Peltigera* (56%). Brine shrimp cytotoxicity is the easiest and simple

method for detection of bioactive compound. In the above study Brine shrimps lethality of the ten lichen extract were shown in (table 4). In acetone extract highest rate of mortality was observed in *Flavopuntelia* (60%) followed by *Flavoparmelia* (40%) and *Peltigera* (30%). For methanol extract highest mortality were recorded for *Flavoparmelia* (53%) followed by *Flavopuntelia* (50%) and *Dermetacarpon*, *Peltigera*, *Usnea* (30%). The result of antitumor activity of different concentration is shown in table (5). In acetone extract of concentration 0.05mg/ml, maximum (40%) antitumor activity was recorded for lichen genera *Flavoparmelia* followed by *Rhizocarpon* (53%), while in methanol extract maximum antitumor activity 47% and 45% for same genera respectively (table 6). Figure 1-4 represent the percentage mortality and tumor inhibition resulting from methanol extract and acetone extract,

**Table 1: Distribution of Different Genera of Lichens collected from Mansehra District Khyber Pakhtoon Khawa, Pakistan.**

| Sr. No. | Lichen genera        | Altitude (m) | Localities    |
|---------|----------------------|--------------|---------------|
| 1       | <i>Dermetacarpon</i> | 1055         | Dader         |
| 2       | <i>Peltigera</i>     | 2580         | Manoor valley |
| 3       | <i>Flavopuntelia</i> | 1050         | Dader         |
| 4       | <i>Collema</i>       | 1030         | Dader         |
| 5       | <i>Flavoparmelia</i> | 2540         | Manoor valley |
| 6       | <i>Lecanora</i>      | 3334         | Lalasar       |
| 7       | <i>Caloplaca</i>     | 3283         | Behisal       |
| 8       | <i>Cladonia</i>      | 2560         | Manoor valley |
| 9       | <i>Rhizocarpon</i>   | 3550         | Lalasar       |
| 10      | <i>Usnea</i>         | 2580         | Manoor valley |

**Table 2: Antibacterial activity of different Lichen extracts**

| Lichen Genera          | Extract  | Zone of inhibitions |                      |                   |                    |                     |                  |
|------------------------|----------|---------------------|----------------------|-------------------|--------------------|---------------------|------------------|
|                        |          | <i>E.coli</i>       | <i>S.epidermidis</i> | <i>S.bacillus</i> | <i>E. faecalis</i> | <i>P.aeruginosa</i> | <i>S. aureus</i> |
| 1 <i>Dermetacarpon</i> | Acetone  | 10.33±0.577         | 8                    | 8.00±1.00         | 8.33±0.577         | 8.33±0.577          | 8.33±0.577       |
|                        | Methanol | 0                   | 7.00±0.23            | 0                 | 9.00±0.577         | 7                   | 7.66±0.577       |
| 2 <i>Peltigera</i>     | Acetone  | 0                   | 0                    | 0                 | 8.66±0.577         | 0                   | 0                |
|                        | Methanol | 0                   | 7                    | 0                 | 0                  | 7                   | 0                |
| 3 <i>Flavopuntelia</i> | Acetone  | 0                   | 9                    | 0                 | 8.00±1.00          | 0                   | 9.00±1.00        |
|                        | Methanol | 8.00±0.47           | 7.00±0.23            | 0                 | 8                  | 8.00±0.47           | 8.66±0.577       |
| 4 <i>Collema</i>       | Acetone  | 0                   | 0                    | 0                 | 0                  | 8.66±0.577          | 8.66±0.577       |
|                        | Methanol | 0                   | 7.00±0.23            | 0                 | 10.66±0.577        | 7±0.23              | 8.00±0.47        |
| 5 <i>Flavoparmelia</i> | Acetone  | 12.66±0.577         | 12.33±1.15           | 9.66±0.577        | 10.33±0.577        | 11.33±0.577         | 11.33±0.577      |
|                        | Methanol | 12.33±0.577         | 7.00±0.23            | 10.33±0.577       | 10                 | 12.33±0.577         | 9.66±0.577       |
| 6 <i>Lecanora</i>      | Acetone  | 10                  | 12.33±0.577          | 10.66±0.577       | 10                 | 7                   | 12.66±0.577      |
|                        | Methanol | 8.33±0.577          | 7.00±0.23            | 9.66±0.577        | 10.66±0.577        | 10                  | 9                |
| 7 <i>Caloplaca</i>     | Acetone  | 8.66±0.577          | 10                   | 0                 | 10.33±0.577        | 10.66±0.577         | 10               |
|                        | Methanol | 10.33±0.577         | 7.00±0.23            | 8.33±0.577        | 10.66±0.577        | 8                   | 8.33±0.577       |
| 8 <i>Cladonia</i>      | Acetone  | 9.66±0.577          | 8.33±0.577           | 8                 | 12.66±0.577        | 9.66±0.577          | 8.33±0.577       |
|                        | Methanol | 0                   | 7.00±0.23            | 0                 | 10                 | 0                   | 0                |
| 9 <i>Rhizocarpon</i>   | Acetone  | 7.66±0.577          | 11.33±0.577          | 10.66±0.577       | 9.33±0.577         | 7.33±0.577          | 8.66±0.577       |
|                        | Methanol | 12.33±0.577         | 7.00±0.23            | 10                | 0                  | 10                  | 7                |
| 10 <i>Usnea</i>        | Acetone  | 8.33±0.577          | 7.66±0.577           | 8.33±0.577        | 7.66±0.577         | 8.33±0.577          | 8                |
|                        | Methanol | 8                   | 7                    | 9                 | 10                 | 10.66±0.577         | 7                |

**Table 3: Antioxidant activity of Methanol extract of different Lichens extract**

| Methanol Extracts |                      | Percent radical scavenging activity |           |           |           |
|-------------------|----------------------|-------------------------------------|-----------|-----------|-----------|
| S. No             | Lichen species       | 1000 µg/ml                          | 500 µg/ml | 250 µg/ml | 125 µg/ml |
| 1                 | <i>Dermetacarpon</i> | 52                                  | 47        | 43        | 36        |
| 2                 | <i>Peltigera</i>     | 58                                  | 56        | 51        | 41        |
| 3                 | <i>Flavopuntelia</i> | 29                                  | 26        | 22        | 20        |
| 4                 | <i>Collema</i>       | 37                                  | 31        | 30        | 25        |
| 5                 | <i>Flavoparmelia</i> | 46                                  | 39        | 33        | 26        |
| 6                 | <i>Lecanora</i>      | 40                                  | 37        | 29        | 26        |
| 7                 | <i>Caloplaca</i>     | 17                                  | 12        | 0         | 0         |
| 8                 | <i>Cladonia</i>      | 66                                  | 54        | 44        | 40        |
| 9                 | <i>Rhizocarpon</i>   | 63                                  | 60        | 56        | 49        |
| 10                | <i>Usnea</i>         | 33                                  | 31        | 26        | 24        |
|                   | Ascorbic acid        | 79                                  | 72        | 65        | 61        |

**Table 4: Antioxidant activity of Acetone extracts of different Lichens extract**

| Acetone extracts |                      | Percent free radical scavenging activity |           |           |           |
|------------------|----------------------|------------------------------------------|-----------|-----------|-----------|
| S. No            | Lichen species       | 1000 µg/ml                               | 500 µg/ml | 250 µg/ml | 125 µg/ml |
| 1                | <i>Dermetacarpon</i> | 48                                       | 46        | 37        | 33        |
| 2                | <i>Peltigera</i>     | 56                                       | 55        | 51        | 38        |
| 3                | <i>Flavopuntelia</i> | 31                                       | 29        | 23        | 21        |
| 4                | <i>Collema</i>       | 26                                       | 19        | 13        | 11        |
| 5                | <i>Flavoparmelia</i> | 51                                       | 48        | 43        | 41        |
| 6                | <i>Lecanora</i>      | 37                                       | 36        | 30        | 28        |
| 7                | <i>Caloplaca</i>     | 29                                       | 20        | 13        | 0         |
| 8                | <i>Cladonia</i>      | 49                                       | 47        | 41        | 33        |
| 9                | <i>Rhizocarpon</i>   | 57                                       | 55        | 46        | 44        |
| 10               | <i>Usnea</i>         | 26                                       | 23        | 16        | 8         |
|                  | Ascorbic acid        | 79                                       | 72        | 65        | 61        |

**Table 5: Showing Cytotoxic activity acetone and methanol extract**

| S.No | Lichen genera        | Acetone Extract concentration |      |       | Methanol extract concentration |      |       |
|------|----------------------|-------------------------------|------|-------|--------------------------------|------|-------|
|      |                      | 0.05                          | 0.01 | 0.001 | 0.05                           | 0.01 | 0.001 |
| 1    | <i>Dermetacarpon</i> | 10%                           | 10%  | 0%    | 30%                            | 30%  | 20%   |
| 2    | <i>Peltigera</i>     | 30%                           | 20%  | 20%   | 30%                            | 20%  | 20%   |
| 3    | <i>Flavopuntelia</i> | 60%                           | 40%  | 30%   | 50%                            | 40%  | 30%   |
| 4    | <i>Collema</i>       | 0%                            | 0%   | 0%    | 0%                             | 0%   | 0%    |
| 5    | <i>Flavoparmelia</i> | 40%                           | 30%  | 26%   | 53%                            | 27%  | 13%   |
| 6    | <i>Lecanora</i>      | 30%                           | 10%  | 0%    | 10%                            | 7%   | 0%    |
| 7    | <i>Caloplaca</i>     | 4%                            | 0%   | 0%    | 10%                            | 0%   | 0%    |
| 8    | <i>Cladonia</i>      | 7%                            | 4%   | 0%    | 10%                            | 7%   | 0%    |
| 9    | <i>Rhizocarpon</i>   | 24%                           | 13%  | 4%    | 27%                            | 17%  | 7%    |
| 10   | <i>Usnea</i>         | 17%                           | 7%   | 0%    | 20%                            | 3%   | 0%    |

**Table 6: Showing Percentage of Tumor inhibition.**

| Percentage of Tumor inhibition |                      |                               |      |       |                                |      |       |
|--------------------------------|----------------------|-------------------------------|------|-------|--------------------------------|------|-------|
| S. NO                          | Lichen species       | Acetone extract concentration |      |       | Methanol extract concentration |      |       |
|                                |                      | 0.05                          | 0.01 | 0.001 | 0.05                           | 0.01 | 0.001 |
| 1                              | <i>Dermetacarpon</i> | 38%                           | 33%  | 26%   | 28%                            | 19%  | 10%   |
| 2                              | <i>Peltigera</i>     | 34%                           | 23%  | 20%   | 30%                            | 16%  | 8%    |
| 3                              | <i>Flavopuntelia</i> | 36%                           | 23%  | 20%   | 38%                            | 30%  | 27%   |
| 4                              | <i>Collema</i>       | 16.66%                        | 13%  | 12%   | 17%                            | 10%  | 0%    |
| 5                              | <i>Flavoparmelia</i> | 40%                           | 38%  | 34%   | 37%                            | 31%  | 15%   |
| 6                              | <i>Lecanora</i>      | 18%                           | 13%  | 0%    | 0%                             | 0%   | 0%    |
| 7                              | <i>Caloplaca</i>     | 0%                            | 0%   | 0%    | 0%                             | 0%   | 0%    |
| 8                              | <i>Cladonia</i>      | 17%                           | 8%   | 0%    | 18%                            | 10%  | 0%    |
| 9                              | <i>Rhizocarpon</i>   | 39%                           | 33%  | 29%   | 37%                            | 25%  | 13%   |
| 10                             | <i>Usnea</i>         | 29%                           | 20%  | 11%   | 28%                            | 20%  | 9%    |




Fig. 1. Percentage mortality rate of Acetone extract




Fig. 2. Percentage of mortality rate of Methanol extract




Fig. 3. Percentage of Tumor inhibition of methanol extract




Fig. 4. Percentage of tumor inhibition of Acetone extract

## DISCUSSION

Our study concluded that all ten genera of lichen had antibacterial activity against most of tested bacterial strains. *S.bacillus* was the most resistant and *S.epidermidis* was the most sensitive bacteria to most of the investigated lichen. According to Aydin and Kinalioglu (2013), lichen *Cetraria aculeate* showed the antimicrobial activity with extracts of acetone, diethyl ether and ethanol. All extracts were found active against bacterial strains of *E.coli*, *Staphylococcus aureus*, *Aeromonas hydrophila*, *Proteus vulgaris*, *Streptococcus faecalis*, *Bacillus cereus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, and *Listeria monocytogenes*.

In the present study, cytotoxicity assay indicated that both methanol and acetone. *Flavopuntelia* were potent against brine shrimp and there mortality rate were 60% and 50% respectively. Babita *et al.* (2012) reported that methanolic extract of lichen *Heterodermia* sp. and *Ramalina* sp. show comparable toxic effect.

The present study also reported that the methanol extract of *Rhizocarpon* have highest free radical scavenging activity in all of the investigated genera. Sharma and Kalikotay 2012 investigated the antioxidant activity of lichen *Parmotrema reticulatum* and *Usnea* sp. They observed 10-35% DPPH free radical scavenging for methanol extract of both lichen species.

In potato disc assay, most of the lichen extract show significant antitumor (inhibition  $\geq 20\%$ ) inhibition activity against *Agrobacterium tumefaciens*. Extract of *Flavoparmelia* show maximum (40%) tumor inhibition while *Collema* show minimum tumor inhibition.

## REFERENCES

- Ali, I (2005). Mapping and documentation of the culture assets of Kaghan valley, Mansehra. UNESCO, Islamabad.
- Aliyu, A. B., M. A. Ibrahim, H. Ibrahim, A. M. Musa, A.Y. Lawal, J.A. Oshaanimi, M. Usman, A. Kadir, A. O. Oyewale (2012). Free radical screening and total antioxidant capacity of methanol extract of *Ethulia conyzoids* growing in Nigeria. Romanian Biot. Lett. 17(4): 7458-7465.
- Aptroot, A. and S. H. Iqbal (2012). Annotated checklist of Lichen of Pakistan, with report of new record. Herzogia. 25(2): 211-229.
- Aydin, S. and K. Kinalioglu (2013). The Investigation of Antibacterial Activities of Ethanol and Methanol Extracts of *Flavoparmelia caperata* (L.) Hale (Parmeliaceae) and *Roccella phycopsis* Ach. (Roccellaceae), lichens Collected from Eastern Blacksea Region, Turkey. J. Appl. Pharma. Sci. 3 (02): 143-147.
- Babita, P., H. D. Bhattarai, D. P. Pandey, J. S. Hur, S. G. Hong, C. Kim and J. H. Yim (2012). Antioxidant, antibacterial and Brine shrimp toxicity test of some Mountainous lichen from Nepal. Biol. Res. 45: 387-391.
- Claudia M. P., E. D. T. Lorena, R. K. Lücking, M. Vizcaya and G. E. Medina (2014). Antioxidant activity, total phenols and flavonoids of lichens from Venezuelan Andes. J. Pharm. Pharmacogn. Res. 2(5): 138-147.
- Coker, P. S., J. Radecke, C. Guy and N. D. Camper (2003). Potato disc tumor induction assay: a

- multiple mode of drug action assay. *Phytomedicine* 10: 133-138.
- Chopta, G. L (1934). *Lichen of the Himalayas. Part. I.* Lahore.
- Culberson, C.F (1972). Improved condition and new data for the identification of lichen product by standardized thin layer chromatographic method. *J. Chromatg.* 72: 113-125.
- Culberson, C.F. and H. Kristinsson (1970). A standard method for the identification of lichen product. *J. of Chromatg.* 46: 85-93.
- Karthikaidevi, G., G. Thirumaran, K. Manivannan, P. Anantharaman, K. Kathiresan and T. Balasubaramanian (2009). Screening of the Antibacterial Properties of Lichen *Rocella belangeriana* (Awasthi) from Pichavaram Mangrove (Rhizophora (Sp.). *Advan in Biol Rese.* 3 (3-4): 127-131.
- Krystle, A., A. Santiago, N. C. Borricano, J. N. Jayne, M. A. Denisse, C. P. Myleen and E. C Thomas (2010). Antibacterial activities of fruticose lichens collected from selected sites in Luzon Island, Philippines. *Phil. Scie. Lett.* 3(2): 18-29.
- Marijana, K. and B. Rankovic (2010). Screening of antimicrobial activity of Lichen species in vitro. *Kragujevac J. Sci.* 32: 65-72.
- Olowa, L. F. and O. M. Muneza (2013). Brine Shrimp Lethality Assay of the Ethanolic Extracts of Three Selected Species of Medicinal Plants from Iligan City, Philippines. *Inter. Res. J. Biol. Sci.* 2(11): 74-77.
- Santiago, K. A. A., E. Sangvichien, K. Boonpragob and C. Tee (2010). Secondary metabolic profiling and antibacterial activities of different species of *Usnea* collected in Northern Philippines. *Mycosphere* Doi 10.5943/mycosphere/4/2/10.
- Sati, S. C. and S. Joshi (2011). Antibacterial Activity of the Himalayan Lichen *Parmotrema nilgherrense* Extracts. *Brit. Microb. Res J.* 1(2): 26-32.
- Shah, N. C (2011). Lichen of commercial Importance in India. *Herb. Tech. Ind.* 10(01): Jan-Feb.
- Sharma, B. C., S. Kalikotay (2012). Screening of antioxidant activity of Lichen *Parmotrema reticulatum* and *Usnea sp* of Darjeeling Hills India. *ISOR.J. Pharm.* 2 (6): 54-60.
- Sharma, B. C., S. Kalikotay and B. Rai (2011). Assessment of antimicrobial activity of extracts of few common Lichen of Darjeeling Hills. *Ind. J. Fund and App. Life Sci.* 2 (1): 120-126.
- Shyam, K., N. Thajuddin and D. K. Upreti (2010). Diversity of Lichen in Kolli hills of Tamil Nadu, India. *Int. J. Biod. Cons.* 3(2): 36-39.
- Smith, A. L (1918). *A Monograph of British the British Lichen.* London.
- Tatjana, M., S. Stamenkovic, V. Cvetkovic, S. Tosic, M. Stankovic, I. Radojevic, O. Stefanovic, L. Comic, D. Dacic, M. Curcic and S. Markovic (2011). Antioxidant, Antimicrobial and Antiproliferative activities of Five Lichen Species. *Int. J. Mol. Sci.* 12: 5428-5448.
- Veranja, K., K. Bombuwela, S. Kathirgamanathar and V. M. Thadhani (2005). Lichen: A chemically important Biota. *J. Nantn. Sci. Foundation Sri Lanka.* 33 (3): 169-186.